WORLD INTELLECTUAL PROPERTY ORGANIZATION



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:		11) International Publication Number: WO 94/00480
C07J 71/00, A61K 31/58	A1	43) International Publication Date: 6 January 1994 (06.01.94)
(21) International Application Number: PCT/US9 (22) International Filing Date: 6 May 1993 (C) (30) Priority data: 07/904,914 26 June 1992 (26.06.92) (60) Parent Application or Grant (63) Related by Continuation US 07/904,9 Filed on 26 June 1992 (26.06.92) (71) Applicant (for all designated States except US): 1 INC. [US/US]; 235 East 42nd Street, New Yor 10017 (US).	06.05.9 U 114 (CI 26.06.9 PFIZE	 (75) Inventors/Applicants (for US only): DENINNO, Michael Paul [US/US]; 9 Hermitage Drive, Gales Ferry, CT 06335 (US). McCARTHY, Peter, Andrew [US/US]; 23: North Anguilla Road, Pawcatuck, CT 06379 (US). (74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 23: East 42nd Street, New York, NY 10017 (US). (81) Designated States: AU, BG, BR, CA, CZ, DE, JP, KR NO, NZ, RO, RU, SK, UA, US, European patent (AT BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM GA, GN, ML, MR, NE, SN, TD, TG).

(54) Title: STEROIDAL GLYCOSIDES FOR TREATING HYPERCHOLESTEROLEMIA

(57) Abstract

Certain steroidal glycosides are useful as hypocholesterolemic agents and antiatherosclerosis agents.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	CN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	1E	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JР	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic	RU	Russian Federation
CF	Central African Republic		of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	K2	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
СМ	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
cs	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	UA	Ukraine
DE	Germany	MG	Madagascar	US	United States of America
DK	Denmark	ML	Mali	UZ	Uzbekistan
ES	Spain	MN	Mongolia	VN	Vict Nam
PI	Finland		•		

-1-

STEROIDAL GLYCOSIDES FOR TREATING HYPERCHOLESTEROLEMIA

5

10

15

20

25

Background of the Invention

This invention relates to steroidal glycosides and methods of using the same, particularly as hypocholesterolemic agents and antiatherosclerosis agents, in mammals.

Many known products possessing hypocholesterolemic activity are cross-linked synthetic polymer derivatives, for example of polystyrene. For example, cross-linked, water-insoluble, bile-acid-binding polystyrene-based resins, e.g., Cholestyramine® agents, have a gritty "mouth-feel", and thus have poor palatability. In addition, these resin beads typically have a low in vivo efficiency. Thus, the effective hypocholesterolemic dose of these materials is excessive, typically 18-24 grams of formulated product per day. Other known polymers having hypocholesterolemic activity include the natural product chitosan and chitosan derivatives as described in European Application pub. no. 0212145. However, the effective hypocholesterolemic dose of these materials is also high.

Other known hypercholesterolemia controlling agents include plant extracts such as "alfalfa saponins". However, these plant extracts are of variable composition and contain significant amounts of nonuseful chemical substances. Due to the variations in composition, it is difficult to set a standard dosage or predict the impurities present. Thus, such extracts are not well suited for use by humans. Furthermore purification of these extracts would be expensive. As an alternative certain synthetically produced, pure "sapogenin-derived" compounds e.g., substances compounded from spirostane, spirostene or sterol-derived compounds depress cholesterol absorption more effectively than alfalfa extracts on a weight basis and thus can be administered in reasonable sized doses. Because the chemical compositions of these substances are known and because they can be synthesized at a high degree of purity, they are suitable for use by any warm-blooded animal, including humans.

10

However, unless administered in massive amounts, pure sapogenins do not significantly inhibit cholesterol's absorption. It is only when compounded with another moiety that sapogenins have the desired effect. Examples of such sapogenin compounds are compounds of tigogenin and diosgenin, particularly glycosides thereof. P.K. Kintia, lu. K. Vasilenko, G.M. Gorianu, V.A. Bobeiko, I.V. Suetina, N.E. Mashchenko, Kim. Pharm. Zh., 1981, 15(9), 55 discloses 3-O-(ß-D-galactopyranosyl)hecogenin and its use as a hypocholesterolemic agent. U.S. Pat. Nos. 4,602,003 and 4,602,005 disclose certain steroidal glycosides, in particular 3-O-(ß-D-glucopyranosyl)tigogenin and 3-O-(ß-D-cellobiosyl)tigogenin and their use for the control of hypercholesterolemia. 3-O-(ß-D-cellobiosyl)tigogenin has superior hypocholesterolemic activity when compared to, for example, cholestyramine.

In addition, certain other steroidal glycosides described below have been 15 published, however these publications do not address hypocholesterolemic activity. "Structural Features of the Antioxidant and fungicidal Activity of Steroid Glycosides", Dimoglo, A. S.; Choban I. N.; Bersuker, I. B.; Kintya, P. K.; Balashova, N. N.; Bioorg. Khim, 11(3), 408-413, 1985 discloses rockogenin B-D-galactopyranoside and tigogenin B-D-lactoside. "Preparation and Properties of Some New Steroid B-D-20 Glucopyranosides, B-D-Glucopyranosiduronic Acids, and Derivatives*. Schneider. J.J.; Carb. Research, 17, 199-207, 1971 discloses tigogenin B-Dglucopyranuronoside. "Sterol Glycoside with Activity as Prostaglandin Synthetase Inhibitor", Pegel, K. H. Walker, H.; United States Patent 4,260,603, April 7, 1981 discloses hecogenin B-D-glucopyranoside. "Hemolytic Properties of Synthetic 25 Glycosides", Segal, R.; Shud, F.; Milo-Goldzweig, I.; J. Pharm, Sci., 67 (11) 1589-1592, 1978 discloses tigogenin ß-D-maltoside, tigogenin ß-L-fucopyranoside, smilagenin ß-maltoside and tigogenin a-L-rhamnoside. "Steroid Glycosides from the Roots of Capsicum Annuum II: The Structure of the Capsicosides, Gutsu, E.V.; Kintya, P.K.; Lazurevskii, G.V.; Khim. Prir. Soedin., (2), 242-246, 1987 discloses 30 tigogenin a-D-arabanopyranoside and tigogenin B-D-galactopyranoside. "Molluscicidal Saponins from Cornus Florida L.", Hostettmann, K.; Hostettmann-Kaldas, M.; Nakanishi, K.; Helv. Chim. Acta, 61, 1990-1995, 1978 discloses smilagenin B-D-galactopyranoside. "Steroidal Saponins from Several Species of Lilifforae Plants*, Yang, C.; Li, K.; Ding, Y.; Yunnan Zhiwu Yanjiu Zengkan, Suppl. 3, 35 13-23, 1990 discloses (25S) - hecogenin cellobioside. "Determination of the

10

20

Absolute Configuration of a Secondary Hydroxy Group in a Chiral Secondary Alcohol Using Glycosidation shifts in Carbon-13 NMR Spectroscopy*, Seo, S.; Tomita, Y.; Tori, K.; Yoshimura, Y.; J. Am. Chem. Soc., 100(11), 3331-3339, 1978 discloses smilagenin β-glucoside and smilagenin φ-glucoside. *Steroid Glycosides from Asparagus Officinalis*, Lazurevskii, G. V.; Goryanu, G. M.; Kintya, P. K.; Dokl. Akad. Nauk. SSSR, 231(6), 1479-81, 1976 discloses sarsasapogenin β-glucoside.

Although the hypocholesterolemic compounds described above make a significant contribution to the art there is a continuing search in this field of art for improved hypocholesterolemic pharmaceuticals.

Summary of the Invention

This invention is directed to steroidal glycosides, particularly spirostanyl glycosides, that are useful as hypocholesterolemic agents and antiatherosclerosis agents. The compounds of this invention have the formula

Formula IA

25

wherein either (A):

$$Q^3$$
 is $C^2 - C^2 + C$

 $R^{1}0$ -alkylene(C_{2} - C_{3})-0 H or

Q4 and Q5 are both methylene;

10 and wherein

R1 is

B-D-glucopyranosyl,

B-D-glucopyranuronosyl,

B-D-2-acetamido-2-deoxy-glucopyranosyl,

β-D-galactopyranosyl,

B-D-fucopyranosyl,

B-L-fucopyranosyl,

B-D-xylopyranosyl,

B-L-xylopyranosyl,

20 α-D-arabanopyranosyl,

α-L-arabanopyranosyl,

a-D-cellobiosyl,

B-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl,

25 β-D-gentiobiosyl,

3-O-B-D-galactopyranosyl-g-D-arabanopyranosyl or

B-D-maltotriosyl;

or (B):

30

Q1, Q4 and Q5 are all methylene;

-5-

$$Q^3$$
 is C^2 , C^2 , C^3 is C^2 , C^3 is C^3 is C^3 is C^4 , C^4

or

R¹0-alkylene(C₂-C₃)-0 H

and wherein

R¹ is

B-D-glucopyranosyl,

B-D-glucopyranuronosyl,

B-D-2-acetamido-2-deoxy-glucopyranosyl,

15 B-D-fucopyranosyl,

B-L-fucopyranosyl,

B-D-xylopyranosyl,

B-L-xylopyranosyl,

α-D-arabanopyranosyl,

a-L-arabanopyranosyl,

20 B-D-cellobiosyl,

B-D-lactosyl,

__ .

B-D-maltosyl,

B-D-gentiobiosyl,

3-O-B-D-galactopyranosyl-a-D-arabanopyranosyl or

25 β-D-maltotriosyl;

or (C):

Q1, Q4 and Q5 are all methylene;

Q² is carbonyl;

30

 $R^{1} = R^{1} = R^{1$

or

5

R¹0-alkylene(C₂-C₃)-0 H

C₂₅ is (R);

and wherein

R1 is

10 B-D-glucopyranuronosyl,

B-D-2-acetamido-2-deoxy-glucopyranosyl,

B-D-fucopyranosyl,

B-L-fucopyranosyl,

B-D-xylopyranosyl,

15 B-L-xylopyranosyi,

a-D-arabanopyranosyl,

α-L-arabanopyranosyl,

B-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl,

20 β-D-gentiobiosyl,

3-O-B-D-galactopyranosyl-a-D-arabanopyranosyl or

B-D-maltotriosyl;

or (D):

Q1, Q2, Q4 and Q5 are each methylene;

and Q^3 is $C^2 + C^2 + C^3 + C^4 + C^4$

30 or

25

$$R^{1}0$$
-alkylene(C_{2} - C_{3})-0 H

and wherein

R1 is

35 β-D-2-acetamido-2-deoxy-glucopyranosyl,

B-D-fucopyranosyl, **B-D-xylopyranosyl**, 5 **B-L-xylopyranosyl**, g-L-arabanopyranosyl, β-D-cellobiosyl, **B-D-gentiobiosyl**, 3-O- β -D-galactopyranosyl- α -D-arabanopyranosyl, or B-maltotriosyl; 10 or (E): Q1, Q2, and Q5 are each methylene; Q4 is carbonyl or 15 C₅ is alpha; C₂₅ is (R); and wherein 20 R1 is B-D-galactopyranosyl, B-D-cellobiosyl, B-D-lactosyl, **B-D-maltosyl** or 25 B-D-maltotriosyl; or (F): Q1, Q2, and Q4 are each methylene; Q5 is carbonyl or

WO 94/00480 PCT/US93/04092

-8-

H OR1 R10 H

O3 is -C- or -C-:

5

C_s is alpha;

C₂₅ is (R); and wherein

R1 is

B-D-galactopyranosyl,

10

B-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl or

B-D-maltotriosyl;

with the proviso that (3B,5a,25R)-3-[(B-D-cellobiosyl)oxy]spirostane

is not included.

A first group of preferred compounds of Formula IA consists of these

compounds wherein Q 1 is carbonyl, $^{-C-}$ or $^{-C-}$, Q 2 is methylene, Q 3 is

20

25

H OR1

, Q⁴ is methylene, Q⁵ is methylene, the C₅ hydrogen is alpha and C₂₅ has

the R configuration. Especially preferred within this group are compounds wherein Q^1 is carbonyl and R^1 is β -D-cellobiosyl, α -D-cellobiosyl, β -D-glucopyranosyl, β -D-galactopyranosyl, β -D-hactosyl, β -D-maltosyl or β -D-maltotriosyl. Also, especially

preferred within this group is a compound wherein Q¹ is C- and R¹ is β-D-cellobiosyl. Another especially preferred compound within this group is a

30

compound wherein Q¹ is -C- and R¹ is β-D-cellobiosyl

10

30

35

A second group of preferred compounds of Formula IA are compounds

wherein Q^1 is methylene, Q^2 is C^2 or C^2 , Q^3 is C^2 , Q^4 is methylene, Q^5 is methylene, the C_5 hydrogen is alpha and C_{25} is (R). Especially preferred within this second group is a compound wherein Q^2 is C^2 and C^3 is C^4 and C^4 is C^4 .

A third group of preferred compounds of Formula IA are compounds wherein

H OH HO H HO H HO H HORL

Q1 is carbonyl, -C- or -C-, Q2 is carbonyl, -C- or -C-, Q3 is -C-,

 Q^4 is methylene, Q^5 is methylene, the C_5 hydrogen is alpha and C_{25} is (R).

compound within this group is a compound wherein Q^1 is Q^2 is carbony and R^1 is B-D-cellobiosyl.

A fourth group of preferred compounds of Formula IA are compounds $H = 0R^{1}$ wherein Q¹ is methylene, Q² is carbonyl, Q³ is $^{-C-}$, Q⁴ is methylene, Q⁵ is

10

methylene, the C_5 hydrogen is alpha and C_{25} is (R). Especially preferred within this fourth group are compounds wherein R^1 is β -D-lactosyl or β -D-cellobiosyl.

C₂₅ is (R). Especially preferred within this fifth group is a compound wherein the C₅ hydrogen is alpha and R¹ is β-D-gentiobiosyl. Another especially preferred compound within this group is a compound wherein the C₅ hydrogen is beta and R¹ is β-D-cellobiosyl.

A sixth group of preferred compounds of Formula IA are compounds wherein

15 Q¹, Q² and Q⁵ are each methylene, Q³ is -C- , Q⁴ is carbonyl, the C₅ hydrogen is alpha and C₂₅ is (R). Especially preferred within this group is a compound wherein R¹ is β-D-cellobiosyl.

A seventh group of preferred compounds of Formula IA are compounds

Yet another aspect of this invention is directed to a method for controlling hypercholesterolemia or atherosclerosis in a mammal by administering to a mammal suffering from hypercholesterolemia or atherosclerosis a hypercholesterolemia or atherosclerosis controlling amount of a Formula I spirostanyl glycoside

35

25

Formula I

wherein

either (A):

5

OH Q1 is methylene, carbonyl,

Q² is methylene, carbonyl,

10

R¹0-alkylene(C₂-C₃)-0 H

or

R¹0-alkylene(C₂-C₃)-0 15

Q⁴ and Q⁵ are both methylene;

and wherein

R¹ is

20 B-D-glucopyranosyl,

B-D-glucopyranuronosyl,

B-D-2-acetamido-2-deoxy-glucopyranosyl,

B-D-galactopyranosyl,

B-D-fucopyranosyl,

25 **B-L-fucopyranosyl**,

B-D-xylopyranosyl,

B-L-xylopyranosyl,

a-D-arabanopyranosyl,

a-L-arabanopyranosyl,

30 a-D-cellobiosyl,

B-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl,

B-D-gentiobiosyl,

3-O-ß-D-galactopyranosyl-o-D-arabanopyranosyl, or B-maltotriosyl; or (B): 5 Q1, Q2, and Q5 are each methylene; 10 Q4 is carbonyl or C₅ is alpha; C₂₅ is (R); and wherein 15 R¹ is B-D-galactopyranosyl, B-D-cellobiosyl, B-D-lactosyl, **B-D-maltosyl** or 20 B-D-maltotriosyl; or (C): Q1, Q2 and Q4 are each methylene; H 0R1 25 Q⁵ is carbonyl or C₅ is alpha; 30 C₂₅ is (R); and wherein; R1 is **B-D-galactopyranosyl**, **B-D-cellobiosyl**, 35

20

25

B-D-lactosyl,

B-D-maltosyl or

5 ß-D-maltotriosyl;

with the proviso that

 $(3\beta,5\alpha,25R)-3-[(\alpha-D-cellobiosyl)oxy]$ spirostane,

(3B.5a.25R)-3-[(B-D-glucopyranosyl)oxy]spirostane,

(3B.5a.25R)-3-[(B-D-cellobiosyl)oxy]spirostane or

(3B,5a,25R)-3-[(B-D-galactopyranosyl)oxy]spirostan-12-one are not included.

A first group of preferred compounds of Formula I are compounds

wherein Q^1 , Q^2 , Q^4 and Q^5 are methylene, C_{25} is (R) and Q^3 is ${}^{-C,-}$. Especially

preferred within this group are compounds wherein the C₅ hydrogen is alpha and R¹ is β-D-glucopyranuronosyl, β-D-maltosyl, β-D-lactosyl, β-D-gentiobiosyl or β-D-galactopyranosyl. Another especially preferred compound within this group is a compound wherein the C₅ hydrogen is beta and R¹ is β-D-cellobiosyl.

A second group of preferred compounds of Formula I are compounds

H OH OH H H OR¹

wherein Q¹ is carbonyl, -C- or -C-, Q² is methylene, Q³ is -C- , Q⁴ and Q⁵

are each methylene, C_{25} is (R) and the C_5 hydrogen is alpha. Especially preferred within this second group are compounds wherein Q^1 is carbonyl and R^1 is β -D-cellobiosyl, σ -D-cellobiosyl, β -D-glucopyranosyl, β -D-galactopyranosyl, β -D-haltotriosyl. Another especially preferred compound within this

second group is a compound wherein Q¹ is G-D-cellobiosyl.

Another especially preferred compound within this second group is a compound

wherein Q¹ is -C- and R¹ is β-D-cellobiosyl.

20

25

30

35

A third group of preferred compounds of Formula I are compounds wherein

H OH HO H H 0 R 1 5 Q 1 is methylene, Q 2 is carbonyl, $^{-C-}$ or $^{-C-}$, Q 3 is $^{-C-}$, Q 4 and Q 5 are

each methylene, C_{25} is (R) and the C_5 hydrogen is alpha. Especially preferred within this third group are compounds wherein Q^2 is carbonyl and R^1 is β -D-cellobiosyl or β -D-lactosyl. Other especially preferred compounds within this third group are

compounds wherein Q² is -C- and R¹ is B-D-cellobiosyl or B-D-galactopyranosyl.

A fourth group of preferred compounds of Formula I are compounds wherein

H OH HO H H OH HO H H OR¹

15 Q¹ is carbonyl, -C- or -C-, Q² is carbonyl, -C- or -C-, Q³ is -C- ,

 Q^4 and Q^5 are each methylene, the C_5 hydrogen is alpha and C_{25} is (R). Especially preferred within this fourth group is a compound wherein Q^1 is carbonyl, Q^2 is carbonyl and R^1 is G-D-cellobiosyl. Especially preferred within this fourth group are

compounds wherein Q^1 is carbonyl, Q^2 is C^{-1} and R^1 is β -D-cellobiosyl or β -D-lactosyl. Another especially preferred compound within this group is a compound wherein Q^1 is C^{-1} , Q^2 is carbonyl, and R^1 is β -D-cellobiosyl. Another especially

preferred compound within this group is a compound wherein Q^1 is C^{-1} , Q^2 is carbonyl and R^1 is B-D-cellobiosyl.

A fifth group of preferred compounds of Formula I are compounds wherein Q^1 , Q^2 and Q^5 are each methylene, Q^3 is Q^4 , Q^4 is carbonyl, the Q^4 hydrogen is alpha and Q^4 is Q^4 . Especially preferred within this group is a compound wherein Q^4 is Q^4 is Q^4 .

A sixth group of preferred compounds of Formula I are compounds wherein

By the Compounds of Formula I are compounds wherein

By the Compounds of Formula I are compounds wherein are compounds wherein Q1, Q2 and Q4 are each methylene, Q3 is C2 and Q5 is carbonyl, the C5 hydrogen is alpha and C25 is (R). Especially preferred within this group is a compound wherein R1 is B-D-cellobiosyl.

This invention is also directed to pharmaceutical compositions for the control of hypercholesterolemia or atherosclerosis in mammals which comprise a compound of the Formula IA and a pharmaceutically acceptable carrier.

Yet another aspect of this invention is directed to a composition comprising a hydrate of a compound of the Formula 1A.

The compounds of Formulas IA and I are herein defined as the single enantiomer having the absolute stereochemistry depicted in Formulas IA and I respectively.

Other features and advantages will be apparent from the specification and claims which describe the invention.

20

15

10

25

Detailed Description of the Invention

Scheme I

15

'Anomerization

20


Deacetylation

25

Scheme II 5 beta peracetylated 10 Mercuric sugar halide Coupling sugar Coupling 15 20 ٧ 25 silylation 30 TMS0-X

۷I

Scheme III



-20-

SCHEME V

5 Oxidation 1) Bromination 2) Elimination ΙX 10 15 20 Epoxidation
 LAH reduction ΧI **25** . 30 H0

H

XII

X۷

15

20

25

30

35

The Formula IA compounds are a subset of the Formula I compounds.

Thus, in the following detailed descriptions of the invention (e.g., how to make the invention, how to use the invention) reference to the Formula I group of compounds, inherently encompasses the Formula IA compounds.

The following description of reaction Schemes I, II & III describe how to make the Formula I compounds wherein Q⁴ and Q⁵ are both methylene.

According to reaction Scheme I, the desired Formula I compounds wherein Q¹, Q² and Q³ are as defined above may be prepared by deacetylating the appropriate alpha peracetylated Formula III compound or beta peracetylated Formula IV compound wherein Q¹ and Q² are as defined above and X is either a bond or alkylene-O-.

Typically the deacetylation is accomplished by combining the Formula III or IV compound with a nucleophilic base such as sodium methoxide or potassium cyanide in a solvent such as methanol, tetrahydrofuran, n-propanol or mixtures thereof at elevated temperatures of about 40°C to about 100°C (typically at reflux) and pressures of 0.5 psi to about 50 psi (typically ambient) for about 0.25 hour to about 2 hours. In addition, for Formula I compounds when the sugar is glucopyranuronosyl, the resultant deacetylated compound is further hydrolyzed by, for example, exposure to sodium hydroxide. Also, where appropriate, those compounds wherein either Q¹ or Q² are carbonyl may be reduced to yield the corresponding alcohols in an alternative process to performing the reduction prior to coupling (described in Reaction Scheme IV and the accompanying text). In an analogous manner, where appropriate, those compounds wherein either Q¹ or Q² are hydroxy may be oxidized to yield the corresponding carbonyl in an alternative process to performing the oxidation prior to coupling.

The desired Formula III compound wherein Q¹ and Q² are as defined above may be prepared by anomerizing the appropriate Formula IV compound wherein Q¹ and Q² are as defined above. The stereochemical terms alpha and beta refer to the configuration of the attachment carbon of the sugar.

Typically the anomerization is performed by treatment with a mineral acid such as hydrobromic acid in an anhydrous aprotic solvent such as methylene chloride at temperatures of 20°C to about 40°C (typically ambient) for at least 24 hours, typically to several days. However, for arabanopyranosyl derivatives the

10

15

20

25

30

35

alpha anomer is obtained directly from the saccharide-steroid coupling described below and the beta anomer from the above process (i.e., the nomenclature reverses).

According to Reaction Scheme II the desired Formula IV compounds wherein Q¹ and Q² are as defined above may be prepared by coupling the appropriate acetylated sugar halide (e.g., bromide) and steroid. More specifically, for those Formula IV compounds where the sugar is other than β-D-maltosyl, β-D-gentiobiosyl or β-D-2-acetamido-2-deoxy-glucopyranosyl, a zinc fluoride promoted coupling of the appropriate Formula V compound (wherein Q¹ and Q² are as defined above and X is either a bond or alkylene-O-) and peracetylated sugar halide is used and for those Formula IV compounds where the sugar is β-D-maltosyl, β-D-gentiobiosyl or β-D-2-acetamido-2-deoxy-glucopyranosyl, a mercuric bromide and mercuric cyanide promoted coupling of the appropriate Formula VI compound (e.g., trimethyl silyl ether of the Formula V compound wherein Q¹ and Q² are as defined above and X is either a bond or alkylene-O-) and peracetylated sugar halide is used.

Generally, the zinc fluoride promoted coupling of the Formula V compound and the peracetylated sugar bromide occurs in a non-protic, anhydrous reaction-inert solvent (e.g., acetonitrile) at a temperature of about 20°C to about 100°C for about 0.5 to about 12 hours. Typically about 0.5 to about 4 equivalents (based on Formula V compound) zinc fluoride is used and about 0.5 to about 3 equivalents acetylated sugar bromide is used. Preferably the coupling is acid catalyzed and it is especially preferred that hydrohalic acid generated during the reaction is used as the acid catalyst. The desired compounds may be prepared at pressures of 0.5 to 50 psi, although typically ambient pressures are used. In a preferred isolation technique the glycosides may be precipitated from the crude filtered reaction mixture (e.g., acetonitrile product solution) by the addition of about 25% to 75% water and the remainder alcohol (e.g., methanol). Precipitation of the product from aqueous methanol/acetonitrile requires less processing than an extractive isolation, and provides a product of greater purity.

Generally, the mercuric bromide and mercuric cyanide promoted coupling of the Formula VI compound and the acetylated sugar bromide is performed in an aprotic, anhydrous solvent such as methylene chloride at a temperature of about 20°C to about 100°C for about 0.5 to about 6 hours. Typically about 0.5 to about 4

15

20

25

30

35

equivalents (based on Formula IV compound) mercuric bromide and mercuric cyanide is used and about 0.5 to about 3 equivalents peracetylated sugar bromide (e.g., β-D-maltosyl, β-D-gentiobiosyl or β-D-2-acetamido-2-deoxy-glucopyranosyl) is used. The desired compounds may be prepared at pressures of 0.5 to 50 psi, although typically ambient pressures are used. Preferably they are isolated as described for the zinc fluoride promoted coupling of the Formula V compound above.

The desired Formula VI compounds wherein Q¹ and Q² are as defined above and X is either a bond or alkylene-O- may be prepared by silylating the appropriate Formula V compound wherein Q¹ and Q² are as defined above and X is either a bond or alkylene-O-.

Generally the Formula V compound, a base such as triethylamine and an activated trialkylsilyl compound (e.g., trimethylsilyl trifluoromethane sulfonate or trimethylsilyl chloride) are reacted in an aprotic, anhydrous solvent such as methylene chloride at a temperature less than about 10°C for about 0.5 hour to about 2 hours.

According to Reaction Scheme III the desired Formula V compounds wherein Q¹ and Q² are as defined above and X is alkylene-O- may be prepared by reducing the appropriate Formula VII compound wherein Q¹ and Q² are as defined above.

Generally the reduction is performed by reaction of the Formula VII compound with lithium aluminum hydride in an anhydrous solvent such as tetrahydrofuran at temperatures of less than about 10°C for about 0.5 hour to about 3 hours.

The desired Formula VII compounds wherein Q¹ and Q² are as defined above may be prepared by coupling the appropriate Formula VIII compound where Q¹ and Q² are as defined above with ethyl diazoacetate in the presence of rhodium acetate dimer. Thus, the Formula VIII compound and ethyl diazoacetate are reacted in an aprotic solvent such as methylene chloride in the presence of rhodium acetate dimer at ambient temperature for about 0.5 hour to about 3 hours.

The starting materials for the above described reaction schemes (e.g., ethyl diazoacetate, peracetylated sugar halides) are readily available or can be easily synthesized by those skilled in the art using conventional methods of organic synthesis.

In addition, as an aid to the preparation of the above steroids, the following paragraphs describe the preparation of the various Formula VIII compounds. Literature references for the preparation of Formula VIII steroid compounds (wherein Q1 is methylene and Q2 and the stereochemistry of the C5 hydrogen and C25 carbon are as defined below) are described in Table I.

Formula VIII Compounds Where Q1 is Methylene and the C₃ Hydroxy Group is Beta

	_
4	\sim
Ł	u
Ţ	v

	C₅ hydrogen	C ₂₅	Q²	Reference
15	а	R	CH ₂	R. E. Marker et. al., J. Am. Chem. Soc.(1943) 65 1199.
	α	R	C=0	Marker et. al., J. Am. Chem. Soc. (1947) 69, 2167.
	α	S	CH,	Goodson & Noller J. Am. Chem. Soc. (1939) 61, 2420.
	σ	S	C=0	Callow & James J. Chem. Soc. (1955) 1671.
20	ß	R	CH₂	Marker et. al., J. Am. Chem. Soc. (1943) 65, 1199.
	В	R	C=0	Marker et. al., J. Am. Chem. Soc. (1947) 69, 2167.
	В	S	CH₂	Marker et. al., J. Am. Chem. Soc. (1943) 65, 1199.
	ß	S	C=0	Kenney & Wall J. Org. Chem. (1957) 22, 468.

25

The following paragraphs describe and/or give literature references for the preparation of the various steroids used as starting materials (i.e., the alternative stereochemistry at the C₃ position and the oxygenation and different epimers at C₁₁ and C12) from the above Formula VIII compounds described in Table I. In general the preparation of the different oxygenated steroids is independent of the stereochemistry at the C₃, C₅ and C₂₅ positions. Thus, once the appropriate stereochemistry at the C₃, C₅ and C₂₅ positions are achieved where Q¹ and Q² are each methylene or where Q1 is methylene and Q2 is carbonyl, the various oxygenated compounds at Q1 and Q2 may be prepared therefrom.

WO 94/00480 PCT/US93/04092

Some of the preparation methods described herein will require protection of remote functionality (i.e., Q¹, Q² and Q³). The need for these protecting groups will vary depending on the nature of the remote functionality and the conditions of the preparation methods. This need is readily determined by one skilled in the art. For a general description of protecting groups and their use, see T.W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1981.

The Formula VIII compounds wherein Q^1 is methylene, Q^2 is either methylene or carbonyl and the C_3 hydroxy is beta may be converted to the corresponding Formula VIII compounds where the C_3 hydroxy is alpha by the following two procedures. These preparative methods may be used independent of the C_{25} stereochemistry.

If Q² is carbonyl, the carbonyl is protected as a ketal (e.g., ethylene ketal), by reacting the steroid with ethylene glycol and an acid catalyst according to the procedure of Engel and Rakhit, Can. J. Chem, <u>40</u>, 2153, 1962. When the C₅ hydrogen is alpha, the C₃ hydroxy group is oxidized to the ketone using pyridinium chloro chromate (PCC) in methylene chloride at ambient conditions. Then the C₃ ketone is reduced with a sterically hindered reducing agent such as K-Selectride[®] reducing agent, at low temperature in tetrahydrofuran to give the C₃ alpha alcohol according to Gondos and Orr, J. Chem. Soc. Chem. Commun. <u>21</u>, 1239, 1982. If appropriate, the Q² protecting group is removed with acid, such as hydrochloric acid, in an appropriate solvent such as acetone.

For those compounds wherein the C_5 hydrogen is beta the same procedures are used as were used when the C_5 hydrogen is alpha except the C_3 ketone is reduced using sodium borohydride in ethanol to furnish the C_3 alpha alcohol.

Reaction Scheme IV illustrates the reaction pathways to achieve the Formula VIII compounds wherein Q¹ and Q² are defined above starting from the Formula VIII compound wherein Q¹ is methylene and Q² is carbonyl.

In general, preparation methods for these compounds may be found in L.F. Fieser and M. Fieser, Steroids, Reinhold Pub. Corp., New York, 1959 and references therein, however, the following descriptive text (which is keyed to Reaction Scheme IV) provides specific guidance.

5

10

15

20

25

15

20

25

30

35

Briefly according to Reaction Scheme IV method 1, the starting material is acetylated and bromininated according to the procedure described in <u>J. Chem. Soc.</u>, 1956, 4344. This intermediate is then reduced with lithium aluminum hydride and treated with silver oxide by a procedure similar to that described in <u>Helv. Act. Chim.</u>, 1953, <u>36</u>, 1241. The resulting ß-11,12-epoxide is opened with trichloroacetic acid, saponified and reduced with zinc and acetic acid using the procedure described in <u>J. Chem. Soc.</u>, 1956, 4330 to give the product shown for method 1.

In method 2, the starting material is selectively acetylated using the procedure described in <u>J. Chem. Soc.</u>, 1956, 430. Using the procedure described in <u>Org. Syn.</u>, 1976, <u>55</u>, 84, the resulting product is oxidized with chromium trioxide and pyridine. Using the procedure described in <u>Synthesis</u>, 1973, 790, the resulting product is saponified with potassium cyanide in water, methanol and THF to give the product shown for method 2.

In method 3, the starting material is converted to the corresponding toluenesulfonylhydrazone which is in turn treated with sodium methoxide using a procedure similar to that described in <u>J. Am. Chem. Soc.</u>, 1954, <u>76</u>, 4013. The resulting 11-ene product is oxidized with osmium tetroxide and N-methylmorpholine-N-oxide according to the procedure describe in <u>Tetrahedron Letters</u>, 1976, 1973 to give the product shown for method 3.

In method 4, the starting material is monobrominated using a procedure described in US Pat. No. 3,178,418. Hydrolysis of this intermediate using the procedure described in <u>J. Chem. Soc.</u>, 1956, 4330 gives the product shown for method 4.

In methods 5 and 6, the starting material is reduced with lithium aluminum hydride according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1954, <u>76</u>, 4013. The two products shown in methods 5 and 6 are separated chromatographically.

In method 7, the starting material is reduced with lithium aluminum hydride according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1951, <u>73</u>, 1777 to give the product shown.

In method 8, the starting material is reduced with lithium and ammonia according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1953, <u>75</u>, 1282 to give the product shown.

In method 9, the starting material is acetylated according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1955, <u>77</u>, 1632 to give a mixture of acetates from which the 3,11-diacetate can be isolated. The unprotected 12-alcohol is then oxidized with chromium trioxide and pyridine according to the procedure described in <u>Org. Syn.</u>, 1976, <u>55</u>, 84. Saponification of the acetates gives the product shown for method 9.

In method 10, the starting material is diacetylated using the procedure described in <u>J. Chem. Soc.</u>, 1956, 4330. The diacetate is reduced with calcium and ammonia using the procedure described in <u>J. Chem. Soc.</u>, 1956, 4334 to give the product shown for method 10.

In method 11, the starting material is reduced with litnium and ammonia according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1953, <u>75</u>, 1282 to give the product shown.

In method 12, the starting material is reduced with lithium aluminum hydride according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1951, <u>73</u>, 1777 to give the product shown.

In method 13, the starting material is selectively protected at the 3-alcohol with t-butyldimethylchlorosilane and imidazole using the procedure described in <u>J. Am. Chem. Soc.</u>, 1972, <u>94</u>, 6190. Using the procedure described in <u>Org. Syn.</u>, 1976, <u>55</u>, 84, the product is oxidized with chromium trioxide and pyridine. The 3-alcohol is then desilylated with hydrofluoric acid in acetonitrile using the procedure described in <u>J. Am. Chem. Soc.</u>, 1972, <u>94</u>, 6190 to give the product shown for method 13.

In method 14, the starting material is selectively protected at the 3-alcohol with t-butyldimethylchlorosilane and imidazole using the procedure described in <u>J. Am. Chem. Soc.</u>, 1972, 94, 6190. The resulting intermediate is reduced with lithium aluminum hydride using the procedure described in <u>J. Am. Chem. Soc.</u>, 1951, <u>73</u>, 1777. The resulting intermediate is selectively acetylated on the 12-alcohol, silylated on the 11-alcohol with trimethylsilyltriflate and 2,6-lutidine using the procedure described in <u>Tetrahedron Letters</u>, 1981, <u>22</u>, 3455, and then deacetylated at the 12-alcohol with lithium aluminum hydride and an aqueous ammonium chloride quench. The 12-alcohol is oxidized with chromium trioxide and pyridine in methylene chloride

30

10

15

20

15

20

25

30

using the procedure described in <u>Org. Syn.</u>, 1976, <u>55</u>, 84, and then desilylated with hydrofluoric acid in acetonitrile using the procedure described in <u>J. Am. Chem. Soc.</u>, 1972, <u>94</u>, 6190 to give the product shown in method 14.

According to Reaction Scheme V the desired Formula IA compounds wherein Q¹, Q², Q³, Q⁴, Q⁵, C₅ and C₂₅ are as described in (E) above (i.e., oxygenated at the Q⁴ position) may be prepared by the following procedures.

The desired Formula X compound can be prepared by the oxidation of tigogenin IX. Generally the oxidation is performed by reaction of tigogenin with pyridinium chlorochromate in a reaction inert solvent such as methylene chloride at 0°C to ambient temperature for about 2 hours to about 10 hours.

The desired Formula XI compound can be prepared by bromination of the Formula X compound followed by an elimination reaction. Typically the bromination is performed by reaction of the Formula X compound with bromine in tetrahydrofuran at a temperature of about -78°C, followed by warming to ambient temperature for about 1 hour to about 3 hours. The elimination reaction is performed by reaction of the brominated product prepared above with lithium bromide and lithium carbonate in a polar, aprotic solvent such as dimethyl formamide at a temperature of about 100°C to about 140°C for about 1 hour to about 4 hours.

The desired Formula XII compound can be prepared by epoxidation of the appropriate Formula XI compound followed by lithium aluminum hydride reduction. Generally the epoxidation is performed by reaction of the Formula XI compound with hydrogen peroxide and sodium hydroxide in a polar, protic solvent such as methanol at ambient temperature for about 2 hours to about 6 hours. The reduction is performed by reaction of the epoxide prepared above with !ithium aluminum hydride in a reaction inert solvent such as tetrahydrofuran at ambient temperature for 2 hours to about 6 hours.

The desired Formula IA compound as described in (E) above wherein Q⁴ contains a hydroxy group may then be prepared from the appropriate Formula XII compound by a zinc fluoride catalyzed coupling followed by deacetylation with

WO 94/00480 PCT/US93/04092

sodium methoxide as described previously. Those compounds wherein Q⁴ is carbonyl may be prepared in an analogous manner, with the addition of oxidation with pyridinium chlorochromate (as described for Formula X compounds) prior to the deacetylation.

According to Reaction Scheme VI, the desired Formula IA compounds wherein Q¹,Q²,Q³,Q⁴Q⁵,C₅ and C₂₅ are as described in (F) above (i.e., oxygenated at the Q⁵ position) may be prepared by the following procedures.

The desired Formula (XIV) compound can be obtained by protection (as denoted by P) of the alcohol function in diosgenin (XIII) followed by hydroboration of the olefin. Typically, the alcohol is protected as an ethoxymethyl ether by reaction of diosgenin with ethoxymethyl chloride and diisopropylethyl amine in an anhydrous solvent such as methylene chloride at ambient temperature for about 2 hours to 6 hours. The hydroboration is performed by reaction of the compound prepared above with borane-tetrahydrofuran complex in a reaction-inert solvent such as tetrahydrofuran at ambient temperature for about 1 hour to about 6 hours.

The desired Formula XV compound can be prepared by oxidation of the appropriate Formula XIV compound followed by removal of the alcohol protecting group. Generally, the oxidation is performed by reaction of the Formula XIV compound with pyridinium chlorochromate in an anhydrous solvent such as methylene chloride at ambient temperature for about 2 hours to about 8 hours. The removal of the alcohol protecting group can be accomplished by reaction of the oxidized product prepared above with concentrated hydrochloric acid in a mixed solvent containing methanol and tetrahydrofuran at a temperature of about 40°C to about 65°C for about 5 minutes to about 1 hour.

The desired Formula IA compound as described in (F) above wherein Q⁵ is carbonyl may then be prepared from the appropriate Formula XV compound by a zinc fluoride catalyzed coupling reaction followed by deacetylation using sodium methoxide as described previously. Those compounds where in Q⁵ contain a hydroxy group may be prepared in an analogous manner with the addition of reduction prior to deacetylation. Typically the reduction is performed by reaction with sodium borohydride in a mixed solvent of ethanol and dichloromethane at ambient temperature for about 1 hour to about 6 hours.

5

10

15

20

25

WO 94/00480 PCT/US93/04092

The compounds of Formula I which have been obtained and have asymmetric carbon atoms can be separated into their diastereomers on the basis of their physical chemical differences by methods known <u>per se.</u>, for example, by chromatography and/or fractional crystallization.

The compounds of this invention where the sugar is β-D-glucopyranuronosyl are acidic and they form base salts. All such base salts are within the scope of this invention and they can be prepared by conventional methods. For example, they can be prepared simply by contacting the acidic and basic entities, usually in a stoichiometric ratio, in either an aqueous, non-aqueous or partially aqueous medium, as appropriate. The salts are recovered either by filtration, by precipitation with a non-solvent followed by filtration, by evaporation of the solvent, or, in the case of aqueous solutions, by lyophilization, as appropriate.

10

15

20

25

30

35

In addition, many of the compounds of this invention may be isolated as hydrates.

The compounds of this invention are potent inhibitors of cholesterol absorption and thus are all adapted to therapeutic use as hypercholesterolemia controlling agents in mammals, particularly humans. Since hypercholesterolemia is closely related to the development of generalized cardiovascular, cerebral vascular or peripheral vascular disorders, secondarily these compounds prevent the development of atherosclerosis particularly arteriosclerosis.

The hypercholesterolemia controlling activity of these compounds may be demonstrated by methods based on standard procedures. For example, the <u>in vivo</u> activity of these compounds in inhibiting intestinal absorption of cholesterol may be determined by the procedure of Melchoir and Harwell (J. Lipid Res., 1985, 26, 306-315).

Activity can be determined by the amount of hypocholesterolemic agent that reduces the cholesterol absorption, relative to the control, in male golden Syrian hamsters. Male golden Syrian hamsters are administered either a cholesterol-free diet (control animals) or a diet supplemented with 1% cholesterol and 0.5% cholic acid for 4 days. The following day the animals are fasted for 18 hours, then administered a 1.5 ml oral bolus of water containing 0.25% methylcellulose, 0.6% Tween 80 and 10% ethanol (control animals) or an oral bolus that contains, in addition, the desired concentration of the compound to be tested. Immediately

10

15

20

25

30

following bolus administration, the animals receive a second 1.5 ml oral bolus of liquid hamster diet containing 1% [3 H] cholesterol (2.0 μ Ci/animal; 210 dpm/nmol) and 0.5% cholic acid, and are fasted for an additional 24 hours. At the end of this second fasting period animals are sacrificed livers are excised, saponified and aliquots are decolorized by addition of hydrogen peroxide, and assessed for radioactivity. Total hepatic radioactivity is calculated based on measured liver weights. The degree of cholesterol absorption is expressed as a percentage of the total radioactivity administered as an oral bolus that is present in the liver 24 hours following bolus administration.

Anti-atherosclerosis effects of the compounds can be determined by the amount of agent that reduces the lipid deposition in the rabbit aorta. Male New Zealand white rabbits are fed a diet containing 0.4% cholesterol and 5% peanut oil for 1 week (meal-fed once a day). After 1 week, the rabbits are dosed daily with the desired concentration of the compound to be tested. After 8.5 weeks, drug treatment is discontinued and the animals are maintained on the cholesterol containing diet for an additional 2 weeks and then switched to a cholesterol free diet for 5 weeks. The animals are sacrificed, and the aortas removed from the thoracic arch to the branch of the iliacs. The aortas are cleaned of adventitia, opened longitudinally and then stained with Sudan IV as described by Holman et al. (Lab. Invet. 1958, 7, 42-47). The percent of the surface area stained is quantitated by densitometry using an Optimas Image Analyzing System (Image Processing Systems). Reduced lipid deposition is indicated by a reduction in the percent surface area stained in the drug treated group in comparison with the control rabbits.

Administration of the compounds of this invention can be via any method which delivers the compounds to the intestinal lumen. These methods include oral routes, intraduodenal routes etc.

The amount of steroidal glycoside administered will, of course, be dependent on the subject being treated, on the severity of the affliction, on the manner of administration and on the judgement of the prescribing physician. However, an effective dosage is in the range of 0.71 to 200 mg/kg/day, preferably 2 to 50 mg/kg/day, most preferably 2 to 7 mg/kg/day. For an average 70 kg human, this

10

15

20

25

would amount to 0.05 to 14 g/day, preferably 0.14 to 3.5 g/day, most preferably 0.14 to 0.5 g/day.

For oral administration, which is preferred, a pharmaceutical composition can take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained release formulations and the like.

Depending on the intended mode of administration, the pharmaceutical compositions may be in the form of solid, semi-solid or liquid dosage forms, such as, for example, tablets, pills, capsules, powders, liquids, suspensions, or the like, preferably in unit dosage forms suitable for single administration of precise dosages. The pharmaceutical compositions will include a conventional pharmaceutical carrier or excipient and a compound according to the invention as an active ingredient. In addition, it may include other medicinal or pharmaceutical agents, carriers, adjuvants, etc.

Pharmaceutical compositions according to the invention may contain 0.1%-95% of the compound, preferably 1%-70%. In any event, the composition or formulation to be administered will contain a quantity of a compound according to the invention in an amount effective to alleviate the signs of the subject being treated, i.e., hypercholesterolemia or atherosclerosis.

For solid pharmaceutical compositions, conventional non-toxic solid carriers include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like.

Liquid pharmaceutically administrable compositions can be prepared by dissolving or dispersing, or otherwise preparing a compound according to this invention and mixing it optionally with a pharmaceutical adjuvant in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol, and the like, to thereby form a solution or suspension.

Methods of preparing various pharmaceutical compositions with a certain amount of active ingredient are known, or will be apparent, to those skilled in this art. For examples, see Remington's Pharmaceutical Sciences., Mack Publishing Company, Easter, Pa., 15th Edition (1975).

WO 94/00480 PCT/US93/04092

-34-

Example 1

(3B,5a,12B,25R)-3-(B-D-galactosyl)oxy]-12-hydroxyspirostane REDUCTION OF KETONES

5

10

15

To a room temperature solution of (3ß,5ø,25R)-3-[(ß-D-galactosyl)oxy]-spirostan-12-one (1.33 g, 2.24 mmol; obtained via deacetylation of (3ß,5ø, 25R)-3-[(tetraacetyl-ß-D-galactosyl)oxy]spirostan-12-one (preparation B2) according to the procedure described in Example 3), ethanol (130 mL) and chloroform (260 mL) was added a solution of sodium borohydride (0.51 g, 13.4 mmol) and ethanol(50 mL). After stirring for 4 hours, methanol (200 mL) was added and stirring was resumed for 2 hours. The reaction mixture was concentrated in vacuo to give 3.38 g of crude product. Recrystallization from a mixture of methanol (80 mL) and water (8 mL), followed by washing with cold methanol (20 mL) and drying gave 1.33 g (quantitative yield) of the title compounds. MS: 595 (M+H).

MP: >200 °C. High resolution FAB MS (m/e): calc. for $C_{33}H_{55}O_9$: 595.3846, found: 595.3861

The title compound was prepared from the appropriate starting material in an analogous manner using the procedure of Example 1.

20

Example 2

(3B,5a,12B,25R)-3-[(B-D-cellobiosyl)oxy]-12-hydroxyspirostane

m.p. M.S.

>200°C 757(M+H) FAB HRMS (m/e): calc. for C₃₉H₆₄O₁₄ Na: 779.4194 779 (M+Na) found: 779.4250

25

Example 3 (38, 5g, 25R)-3-[(β-D-cellobiosyl)oxy]spirostan-11-one

DEACETYLATION

A mixture of (36,5a,25R)-3-[(heptaacetyl-6-D-cellobiosyl)oxy]spirostan-11-one (6.57 g, 6.26 mmol), sodium methoxide (68 mg, 1.25 mmol), methanol (35 mL) and tetrahydrofuran (75 mL) was heated to reflux for 1 hour, followed by stirring at room temperature for 12 hours. A white precipitate formed within 30 minutes. The final

10

15

20

25

30

follows:

suspension was concentrated <u>in vacuo</u> to give 6.0 g of crude product. This material was purified by flash chromatography (eluent: chloroform followed by 8:2 chloroform:methanol) to give 2.71 g (57% yield) of the title compound.

¹H NMR (DMSO-d₆) δ: 5.22 (d, J= 5Hz, 1H); 5.00 (m, 3H); 4.64 (s, 1H); 4.58 (t, J= 5 Hz, 1H); 4.54 (t, J= 6Hz, 1H); 4.34 (q, J= 8 Hz, 1H); 4.27 (d, J= 8Hz, 1H); 4.23 (d, J= 8 Hz, 1H); 3.68 - 2.94 (m, 15 H); 2.34 (m, 2H); 2.08 - 0.81 (m, 23H); 0.92 (s, 3H); 0.86 (d, J= 7 Hz, 3H); 0.72 (d, J= 6 Hz, 3H); 0.59 (s, 3H). DEPT ¹³C NMR (DMSO-d₆) δ: 210.4 (s), 108.8 (s), 103.6 (d), 100.6 (d), 81.1 (d), 80.6 (d), 77.2 (d), 76.9 (d), 76.5 (d), 75.5 (d), 75.1 (d), 73.7 (d), 73.6 (d), 70.5 (d), 66.4 (t), 63.5 (d), 61.5 (t), 60.9 (t), 50.5 (d), 57.1 (t), 54.7 (d), 44.3 (s), 44.1 (d), 41.7 (d), 36.8 (d), 35.6 (t), 35.2 (s), 34.0 (t), 32.6 (t), 31.3 (s), 30.2 (d), 29.2 (t), 28.9 (t), 28.2 (t), 17.5 (q), 17.3 (q), 14.8 (q), 12.3 (q). IR (KBr): 3407 (s), 1700 (m) cm⁻¹. High resolution FAB MS (m/e): calculated for $C_{39}H_{62}O_{14}Na$ 777.4037, found 777.4108. Analysis: calc. for $C_{39}H_{62}O_{14}$ 2H₂O, C 59.22 H 8.41; found C 59.48, H 8.48. MP: >300°C. A monohydrate crystalline form of the above titled product was prepared as

A mixture of 20 g of the crude product prepared according to the above procedure, 600 ml of n-propanol, and 400 ml of water was stirred and heated to reflux. To the resulting solution was charged 2.0 g of diatomaceous earth. While still at reflux the insolubles were removed by filtration. The filtrate was atmospherically distilled to a total volume of 600 ml and cooled to ambient temperature. The resulting suspension was granulated for one hour and the product was collected by filtration. The undried recrystallized cake from the above recrystallization was suspended in 500 ml of methanol. This suspension was heated to reflux for 16 hours, cooled to ambient temperature, granulated for 48 hours, and isolated by filtration. Vacuum drying yielded 16.1 g (81% recovery) of the crystal monohydrate of the Example 3 titled compound.

Examples 4-47

The following compounds were prepared from the appropriate starting material in an analogous manner using the above procedures.

_		m.p.	M.S.	formula	elemental analysis
5	====	===			
	4.)	(3B,5a,25R)-3 >200°C	- <u>{(ß-D-cellobios</u> 755 (M+H) 777 (M+Na)	syl)oxy]spirosta C ₃₉ H ₆₂ O ₁₄ -2H ₂ O	n-12-one calc. C 59.22; H 8.41 found C 59.54; H 8.64
	5.)	(3a,5a,25R)-3	-[(B-D-cellobios	syl)oxy]spirosta	<u>ne</u>
10		>300°C	741 (M+H)	C ₃₉ H ₆₄ O ₁₃ ·0.5H ₂ O	calc. C 62.46; H 8.74 found C 62.31; H 8.36
	6.)	(3B,5B,25R)-3	-[(B-D-cellobios	syl)oxylspirosta	ne
45		>300°C	741 (M+H)	C ₃₇ H ₆₄ O ₁₃ ⋅H ₂ O	calc. C 61.72; H 8.77 found C 61.76; H 9.04
15	7.)	(3B,5a,25R)-3	-[(ß-D-glucuror	nosyl)oxy]spiro	stane
	>2	200°C 615 (M 637 (M		IMS (m/e): calc	t. for C ₃₃ H ₅₁ O ₉ Na₂ 637.3278 found 637.3329
20	8.)	(3β,5α,25R)-3	-[(ß-D-glucosy	l)oxy]spirostan	-12-one
		>200°C	593 (M+H) 615 (M+Na)	C ₃₃ H ₅₂ O ₉ ⋅H ₂ O	calc. C 64.89; H 8.91 found C 64.46; H 8.62
	9.)	(3B,5a,25R)-3	3-[(B-D-galactos	syl)oxy]spiroste	<u>ın-11-one</u>
25		>200°C	593 (M+H) 615 (M+Na)	C ₃₃ H ₅₂ O ₉ ⋅0.75 H ₂ O	calc. C 65.38; H 8.89 found C 65.34; H 8.63
	10.)	(38,58,258)-3	3-[(B-D-cellobio	syl)oxy]spirost	ane
		>250°C	741 (M+H)	C ₃₉ H ₆₄ O ₁₃ ·1.5 H ₂ O	calc. C 60.99; H 8.85 found C 60.69; H 8.90
30	11.)	<u>(3β,5α,25R)-</u>	3-([(ß-D-cellobi	osyl)oxy]ethox	y)spirostane
		>250°C	785(M+H)	C ₄₁ H ₆₈ O ₁₄ ·1.5 H ₂ O	calc. C 60.65; H 8.81 found C 60.53; H 8.97

5	===:	m.p.	M.S. =======	formula ======	elemental analysis		
	12.)	(38,5a,25R)-3	-([(B-D-galacto	pyranosyi)oxy]	ethoxy)spirostane		
		225°C (dec)	623(M+H)	C ₃₅ H ₅₈ O ₀ ·1.25 H ₂ O	calc. C 65.14; H 9.45 found C 65.39; H 9.61		
10	13.)	(3β,5ø,25R)-3	-[(ß-D-maltosyl)oxy]spirostane	<u>e</u>		
		230°C (dec)	741(M+H)	C ₃₉ H ₆₄ O ₁₃ ·2 H₂O	calc. C 60.30; H 8.82 found C 60.64; H 8.84		
	14.)	(3B,5ø,25R)-3	-[(ß-D-lactosyl)	oxy)spirostane	!		
15		>260°C	741(M+H)	C ₃₉ H ₆₄ O ₁₃	calc. C 63.22; H 8.71 found C 6296; H 8.65		
	15.)	(3ß,5a,25R)-3-[(ß-D-lactosyl)oxy]spirostan-12-one					
		>260°C	755(M+H) -	C ₃₉ H ₆₂ O ₁₄ -0.5 H₂O	calc. C 61.31; H 8.31 found C 61.02; H 8.45		
20	16.)	(38,5ø,25R)-3 spirostane	- <u>[(ß-D-2-acetar</u>	<u>nido-2-deoxygl</u>	ucopyranosyl)oxyl-		
		210-212°C	620(M+H)	C ₃₉ H ₆₂ O ₁₄	calc. C 65.91; H 9.32; N 2.20		
				-1.0 H ₂ O	· · = ·		
25	17.)	(3B,5a,25R)-3	B-[(B-D-gentiob	iosyl)oxy]spiros			
		265°C (dec)	741(M+H)	C ₃₉ H ₆₄ O ₁₃ ·1.0 H ₂ O	calc. C 61.72; H 8.77 found C 61.71; H 8.96		
	18.)	(3B,5a,25R)-3	3-[(<i>a</i> -L-arabanc	pyranosyl)oxy	spirostane		
30		>200°C	549(M+H)	C ₃₂ H ₅₂ O ₇ ⋅0.75 H ₂ O	calc. C 68.36; H 9.59 found C 68.30; H 9.64		
	19.)	(3B,5\alpha,25R)-3	3-[(a-D-araban	opyranosyl)oxy	<u>]spirostane</u>		
		>200°C	549(M+H)	C ₃₂ H ₅₂ O ₇ ·2.0 H ₂ O	calc. C 65.73; H 9.65 found C 65.54; H 9.25		

-38-

5		m.p.	M.S.	formula	elemental analysis		
5	20.)	(3ß,5ø,25R)-3-	[(ß-L-xylopyrar	nosyl)oxy]spiro	<u>stane</u>		
	·	>230°C	549(M+H)	C ₃₂ H ₅₂ O ₇ -0.5 H ₂ O	calc. C 68.91; H 9.58 found C 68.52; H 9.36		
10	21.)	(3B,5a,25R)-3-	·[(ß-L-fucopyra	nosyl)oxy]spiro	stane		
		>230°C	561(M+H)		calc. C 68.25; H 9.72 found C 68.62; H 9.53		
	22.)	(3β,5α,25R)-3-	-[(ß-D-xylopran	osyl)oxy]spiros	stane		
15		>220°C	549(M+H)	C ₃₂ H ₅₂ O ₇ ·1.75 H ₂ O	calc. C 66.23; H 9.64 found C 66.32; H 9.31		
	23.)	(3B,5a,25R)-3-[(B-D-fucopyranosyl)oxy]spirostane					
		>220°C	563(M+H)	C ₃₃ H ₅₄ O ₇ -0.5 H ₂ O	calc. C 69.32; H 9.69 found C 69.32; H 9.78		
20	24.) (3β,5α,25R)-3-[(β-D-galactopyranosyl)oxy]spirostane						
		>200°C	579(M+H)	C ₃₃ H ₅₄ O ₈ -1.25 H ₂ O	calc. C 66.12; H 9.47 found C 66.46; H 9.26		
05	25.)	(38,5\alpha,25R)-3-[(3-O-\beta-D-galactopyranosyl)-\alpha-D-arabanopyranosyl)oxy]spirostane					
25		>200°C	579(M+H)		calc. C 62.64; H 8.78 found C 62.93; H 8.66		
	26.)	(3B,5a,25S)-3	-[(ß-D-galactop	oyranosyl)oxy]s	pirostane		
30		>200°C	579(M+H)	C ₃₃ H ₅₄ O ₈ ·2 H₂O	calc. C 64.51; H 9.44 found C 64.69; H 9.41		
	27.)	(3β,5α,25R)-3	-[(a-D-cellobio	syl)oxy]spirosta	<u>an-11-one</u>		
290-292°C 777(M+Na) FAB HRMS (m/e): calc. for $C_{39}H_{62}O_{14}$ 755 found 755.4163							

WO 94/00480 PCT/US93/04092

-39-

5		m.p.	м.s.	formula	elemental analysis			
	#2=2==================================							
	28.)	(3ß,5ø,12ß,25 11-one	iR)-3-[(ß-D-cello	obiosyl)oxy]-12	-hydroxyspirostane-			
10		>200°C	771(M+H) 793(M+Na)	C ₃₉ H ₆₂ O ₁₅ H ₂ O	calc. C 60.76; H 8.10 found C 61.76; H 8.88			
	29.)	(3B,5a,11a,25	5R)-3- <u>[</u> (β-D-cello	obiosyl)oxy]-11	-hydroxyspirostane			
	>210	°C 779(M+N	a) FAB HRMS	(m/e): calc. fo	r C ₃₉ H ₆₄ O ₁₄ Na 779.4194 found 779.4138			
45	30.)	(3β,5σ,11β,25	5R)-3-[(ß-D-cell	obiosyl)oxy]-11	-hydroxyspirostane			
15		>210°C	779(M+Na)	C₃₀H₀₄O₁₄ H₂O	calc. C 60.45; H 8.28 found C 60.41; H 8.58			
	31.)	(3ß,5a,25R)-3	3-[(B-D-glucopy	ranosyl)oxy]sp	irostan-11-one			
20		293-295°C	593(M+H)	C ₃₃ H ₅₂ O ₉ -2 H ₂ O	calc. C 64.89; H 8.91 found C 64.48; H 8.85			
	32.)	(38,50,118,1) di(hydroxy)s		-cellobiosyl)ox	<u>y]-11,12-</u>			
		>230°C	773(M+H)	C ₃₉ H ₆₄ O ₁₅ -2 H ₂ O	calc. C 57.91; H 8.47 found C 57.87; H 8.41			
25 .	33.)	(38,5a,11a,1 di(hydroxy)s		-cellobiosyl)ox	y <u>]-11,12-</u>			
		>230°C	773(M+H)	C ₃₉ H ₆₄ O ₁₅ ·1H ₂ O	calc C 59.22; H 8.41 found C 59.27; H 8.32			
	34.)	(38,5 a ,12 a ,2	5R)-3-[(ß-D-cel	lobiosyl)oxyl-1:	2-hydroxyspirostane			
30		>230°C	757(M+Na)	C₃₃H₅₄O₁₄ ∙3H₂O	calc. C 57.76; H 8.70 found C 57.56; H 8.61			

_		m.p.	M.S.	formula	elemental analysis		
5	======	======	======	======	: = = = = = = = · · · · · · · · · · · ·		
	35.)	(3β,5σ,25R)-3-[(β-D-lactosyl)oxy]spirostan-11-one					
		>270°C	755(M+H)	C ₃₉ H ₆₂ O ₁₄	calc. C 62.05; H 8.28 found C 61.88; H 8.14		
10	36.)	(38,5a,11a,12 di(hydroxy)sr	2 <u>a,25R)-3-[(ß-D</u> birostane	-cellobiosyl)ox	y <u>]-11,12-</u>		
	287-2	88°C 795(M+	H) FAB HRMS	6 (m/e): calc. t	ior C ₃₉ H ₆₄ O ₁₅ Na 795.4143 found 795.4164		
15	37.)	(3B,5ø,11ø,25 12-one	5 <u>R)-3-[(ß-D-cell</u>	obiosyl)oxy]-11	-hydroxyspirostan-		
15		>300°C	771(M+H) 793(M+Na)	C ₃₉ H ₆₂ O ₁₅ -2.5H ₂ O			
	38.)	(3B,5a,25R)-3	B-[(B-D-cellobio	syl)oxy]spirost	an-11,12-dione		
20		244-246°C	769(M+H) 791(M+Na)	C ₃₈ H ₆₀ O ₁₅ 4H ₂ O	calc. C 55.70; H 8.15 found C 55.93; H 7.99		
	39.)	(3β,5α,11β,12α,25R)-3-[(β-D-cellobiosyl)oxy]-11,12- di(hydroxy)spirostane					
	>228	-229°C 773(M 795(M+Na)	I+H) FAB HRM	MS (m/e): calc.	for C ₃₉ H ₆₄ O ₁₅ Na 795.41429 found 795.4164		
25	40.)	(38,5a,12a, 2 11-one	25R)-3-[(ß-D-ce	llobiosyl)oxy]-1	2-hydroxyspirostan-		
		>230°C	771 (M+H) 793 (M+Na)	C ₃₉ H ₆₂ O ₁₅ 3H ₂ O	calc. C 56.78; H 8.31 found C 56.97; H 7.80		
30	41.)	(3ß,5ø,12ß, 2	25R)-3-[(ß-D-lac	tosyl)oxy]-12-h	ydroxyspirostan-11-one		
		>275°C	771(M+H)	C ₃₉ H ₆₂ O ₁₅ -0.5H ₂ O	calc. C 60.06; H 8.14 found C 59.92; H 7.89		

35

Example# Compound Name

5	=======	m.p. =======	M.S.	formula	elemental analysis
	42.)	(3B,5a,25R)-3	-[(B-D-maltotric	osyl)oxy]spiros	tan-11-one
		>230°C	939(M+Na)	C ₃₉ H ₆₂ O ₁₅ 4.5H ₂ O	calc. C 54.15; H 8.17 found C 54.08; H 7.84
10	43.)	(3β,5α,25R)-3	-[(ß-D-maltosy	l)oxy]spirostan	<u>-11-one</u>
		>280°C	755(M+H)	C ₃₉ H ₆₂ O ₁₄ -2H ₂ O	calc. C 59.22; H 8.41 found C 59.38; H 8.13
	44.)	(1a,3ß,5a,25F	R)-3-[(ß-D-cello	biosyi)oxy]-1-h	ydroxyspirostane
15		>280°C	757(M+H)	30 07 17	calc. C 60.45; H 8.58 found C 60.30; H 8.21
	45.)	(3B,5ø,25R)-3	3-[(ß-D-cellobio	syl)oxy]spirost	an-1-one
20		>250°C	755(M+H)		calc. C 60.32; H 8.36 found C 60.24; H 8.13
	46.)	(3β,5α,25R)-3	3-[(ß-D-cellobio	syl)oxy]spirost	an-6-one
		>200°C 754	4(M+H) FAB H	HMRS (m/e): c	ealc. for C ₃₉ H ₆₂ O ₁₄ 755.4286 found 755.4219
25	47.)	(3B,5a,6a,25	R)-3-[(ß-D-cello	biosyl)oxy]-6-h	ydroxyspirostane
		>250°C	756(M+H)	C ₃₉ H ₆₄ O ₁₄ -2H ₂ O	calc. C 59.12; H 8.65 found C 59.23; H 8.46

Example 48

$\underline{(3\beta,5\alpha,11\beta,25R)-3-[(\beta-D-cellobiosyl)oxy]-11-hydroxyspirostan-12-one}$

DEACETYLATION

Based on the procedure described in <u>Synthesis</u>, 1973, 790, (3ß,5\alpha,11\beta,25R)-3-[(heptaacetyl-\beta-D-cellobiosyl)oxy]spirostan-11-ol-12-one (240 mg, 0.225 mmol) was dissolved in methanol (20 mL) and tetrahydrofuran (10 mL). To this solution was added a solution of potassium cyanide (146 mL, 2.25 mmol) in water (0.1 mL) and

methanol (5 mL). The resulting mixture was heated to 80 ° C for four hours. After cooling, the mixture was concentrated to dryness and purified by flash chromatography (9:1 chloroform:methanol eluent) to give the title compound. Mp 245-247 °C. MS (m/e): 771 (P+1), 793 (P+Na). Analysis: calc for C₃₉H₆₂O₁₅ 3H₂O, C 56.70, H 8.31; found, C 56.97, H 7.80.

Preparation A1

(38,5\alpha,25R)-3-[(Heptaacetyl-\alpha-D-cellobiosyl)oxy]spirostan-11-one ANOMERIZATION

Hydrobromic acid (30% in acetic acid, 1.2 mL) was added to a room temperature solution of (36, 5a, 25R)-3-[(heptaacetyl-6-D-cellobiosyl)-oxy]spirostan-11-one (2.0 g) in methylene chloride (35 mL) and the resulting mixture was stirred at room temperature for 94 hours. The reaction was quenched by slow addition of saturated aqueous sodium bicarbonate (20 mL). The organic layer was separated, dried over magnesium sulfate, and dried in vacuo to give 1.637 g of a black solid. Purification by repeated flash chromatography (1:1 hexane:ethyl acetate eluent) provided 651 mg (33% yield) of the title compound.

20

25

15

10

¹H NMR (CDCl₂) δ : 5.41 (t, J = 10 Hz, 1H); 509 (complex, 3H); 4.91 (t, J = 8 Hz, 1H); 4.69 (dd, J = 4 & 10 Hz, 1H); 4.49 (complex, 3H);4.36 (dd, J = 4 & 13 Hz, 1H); 3.99 (m, 3H); 3.67 (m, 2H); 3.40 (m, 3H);2.45 (m, 1H); 2.22 (s, 2H); 2.11 (s, 3H); 2.07 (s, 3H); 2.03 (s, 6H); 2.00 (s, 3H); 1.99 (s, 3H); 1.97 (s, 3H); 2.00 - 0.80 (m, 22H); 1.02 (s, 3H); 0.92 (d, J = 7 Hz, 3H); 0.77 (d, J = 7 Hz, 3H); 0.69 (s, 3H). DEPT ¹³C NMR (CDCl₃) δ : 210.0 (s), 170.5 (s), 170.3 (s), 170.2 (s), 169.6 (s), 169.3 (s), 169.1 (s), 109.2 (s), 100.9 (d), 94.3 (d), 80.6 (d), 78.0 (d), 77.0 (d), 73.1 (d), 71.9 (d), 71.8 (d), 71.2 (d), 69.6 (d), 68.1 (d), 67.8 (d), 66.9 (d), 64.4 (d), 62.0 (t), 61.5 (t), 60.7 (d), 57.6 (t), 55.7 (d), 45.0 (d), 44.3 (s), 41.8 (d), 36.9 (d), 35.5 (t), 35.4 (t), 35.1 (s), 32.7 (t), 31.3 (t), 31.2 (t), 30.2 (d), 28.7 (t), 28.0 (t), 27.4 (t), 20.9 (q), 20.7 (q), 20.6 (q), 20.5 (q), 17.1 (q), 17.0 (q), 14.2 (q), 12.1 (q). IR (Kbr): 1751 (s), 1706 (m) cm⁻¹. MS (m/e): 1049 (M+H), 1071 (M + Na). Analysis: calc. for C₅₃H₇₆O₂₁·H₂O, C 59.65 H 7.37; found C 59.66 H 7.00. MP: 248-249°C.

30

10

15

-43-

Preparation B1

(3ß,5g,25R)-3-[(Heptaacetyl-ß-D-cellobiosyl)oxy]spirostane-11-one ZINC FLUORIDE PROMOTED COUPLING OF FREE SPIROSTANE

A suspension of (3ß,5ø,25R)-3-hydroxyspirostan-11-one (3.0 g, 6.97 mmol) and anhydrous zinc fluoride (2.88 g, 27.9 mmol) in dry acetonitrile (175 mL) was dried by removal of 75 mL of acetonitrile by distillation. The suspension was allowed to cool, heptaacetyl-ß-D-cellobiosyl bromide (9.75 g, 13.9 mmol) was added and the resulting suspension was heated to 65°C for 3 hours. After cooling to room temperature, methylene chloride (150 mL) was added, the suspension was stirred for 10 minutes and filtered. The filtrate was concentrated in vacuo to give 10 g of crude product. This material was dissolved in 8:2 chloroform:methanol, preadsorbed on silica gel and purified by flash chromatography (eluent: 1:1 ethyl acetate:hexane followed by pure ethyl acetate) to give 6.81 g (93% yield) of the title material.

¹H NMR (CDCl₂) δ : 5:11 (complex, 2 H); 5.04 (t, J = 9 Hz, 1H); 4.90 (t, J = 9 Hz, 1H); 4.83 (t, J = 8 Hz, 1H); 4.49 (complex, 4H); 4.34 (dd,J = 4.5 & 12.5 Hz, 1H); 4.04 (t, J = 13 Hz, 1H); 4.03 (t, J = 11 Hz, 1H); 3.72 (t, J = 9.5 Hz, 1H); 3.65 (m, 1H); 3.56 (m, 1H); 3.45 (m, 1H); 2.47 (m, 1H); 2.22 (s, 2H); 2.08 (s, 3H); 2.06 (s, 3H); 2.00 (s, 6H); 1.99 (s, 6H); 1.96 (s, 3H); 2.00 - 1.00 (m, 22H); 0.98 (s, 3H); 0.92 (d, J = 7 Hz, 3H); 0.77 (d, J = 7 Hz, 3H); 0.68 (s, 3H). DEPT ¹³CNMR (CDCl₃) δ : 209.9 (s), 170.5 (s), 170.3 (s), 170.2 (s), 169.9 (s), 169.8 (s), 169.5 (s), 169.3 (s), 169.0 (s), 109.2 (s), 100.8 (d), 99.4 (d), 90.0 (s), 80.6 (d), 79.4 (d), 76.6 (d), 75.3 (s), 72.9 (d), 72.6 (d), 72.5 (d), 71.9 (d), 71.8 (d), 71.6 (d), 67.8 (s), 66.9 (t), 64.4 (d), 62.1 (t), 61.5 (t), 60.8 (s), 60.7 (d), 57.6 (t), 55.7 (d), 44.8 (d), 44.3 (s), 41.8 (d), 36.9 (d), 35.6 (t), 35.2 (s), 34.1 (t), 32.7 (t), 31.3 (t), 31.2 (t), 30.2 (d), 29.0 (t), 28.7 (t), 28.0 (t), 20.9 (q), 20.7 (q), 20.6 (q), 20.5 (q), 20.5 (q), 17.1 (q), 17.0 (q), 14.2 (q), 12.0 (q). IR (KBr): 1756 (s), 1706 (m) cm⁻¹ 1. MS (m/e): 1049 (M+H). Analysis: calc. for C₅₃H₇₆O₂₁·H₂O, C

59.65, H 7.37; found C 59.86, H 7.25. MP: 210-212°C.

20

25

	In an analogous manner the following compounds, Preparation B2-B31, were
	prepared from the appropriate starting material using the above general procedure.
5	Preparation B2
	(38,5a,25R)-3-[(tetraacetyl-6-D-galactosyl)oxy]spirostane-12-one
	Preparation B3
	(3a,5a,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]spirostane
	Preparation B4
10	(3B,5B,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]spirostane
	Preparation B5
	(3B,5a,25R)-3-[(triacetyl-B-D-glucuronosyl)oxy]spirostane methyl ester
	Preparation B6
	(3B,5a,25R)-3-[(tetraacetyl-B-D-glucopyranosyl)oxy]spirostane-12-one
15	Preparation B7
	(33,5a,25R)-3-[(tetraacetyl-3-D-galactopyranosyl)oxy]spirostane-11-one
	Preparation B8
	(3B,5a,25S)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]spirostane
	Preparation B9
20	(3B,5g,25R)-3-([(heptaacetyl-B-D-cellobiosyl)oxy]ethoxy)spirostane
	Preparation B10
	(38,5a,25R)-3-([(tetraacetyl-8-D-galactopyranosyl)oxy]ethoxy)spirostane
	Preparation B11
	(3B,5a,25R)-3-[(heptaacetyl-B-D-lactosyl)oxy]spirostane
25	Preparation B12
	(3B,5g,25R)-3-[(heptaacetyl-B-D-lactosyl)oxy]spirostane-12-one
	Preparation B13
	(3ß,5ø,25R)-3-[(triacetyl-ø-L-arabanopyranosyl)oxy]spirostane
	Preparation B14
30	(3ß,5a,25R)-3-[(triacetyl-a-D-arabanopyranosyl)oxy]spirostane
	Preparation B15
	(38,5a,25R)-3-[(triacetyl-ß-L-xylopyranosyl)oxy]spirostane
	Preparation B16

(3B,5a,25R)-3-[(triacetyl-B-L-fucopyranosyl)oxy]spirostane

-45-

	Preparation B17
5	(3B,5g,25R)-3-[(triacetyl-B-D-xylopyranosyl)oxylspirostane
	Preparation B18
	(3β,5σ,25R)-3-[(triacetyl-β-D-fucopyranosyl)oxy]spirostane
	Preparation B19
	(38,5a,25R)-3-[(tetraacetyl-B-D-galactopyranosyl)oxy]spirostane
	Preparation B20
10	(3B,5a,25R)-3-[(hexaacetyl-3-0-B-D-galactopyranosyl-a-D-arabanopyranosyl)oxy]-spirostane
	Preparation B21
	(3B,5a,25S)-3-[(tetraacetyl-B-D-galactopyranosyl)oxy]spirostane
	Preparation B22
15	(3B,5a,12B,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]-12-hydroxyspirostan-11-one
	Preparation B23
	(3B,5a,11a,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]-11-hydroxyspirostane
20	Preparation B24
	(3B,5g,11B,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]-11-hydroxyspirostane
	Preparation B25
	(3B,5a,25R)-3-[(tetraacetyl-B-D-glucopyranosyl)oxy]spirostan-11-one
25	Preparation B26
	(38,5\alpha,11\beta,12\beta,25\beta)-3-[(heptaacetyi-\beta-D-cellobiosyl)oxy]-11,12- di(hydroxy)spirostane
	Preparation B27
30	(38,5ø,11ø,128,25R)-3-[(heptaacetyl-8-D-cellobiosyl)oxy]-11,12- di(hydroxy)spirostane
	Preparation B28
	(38,5g,12g,25R)-3-[heptaacetyl-8-D-cellobiosyl)oxy]- 12-hydroxyspirostane

WO 94/00480 PCT/US93/04092

-46-

	Preparation B29
5	(3β,5σ,25R)-3-[(heptaacetyl-β-D-lactosyl)oxy]spirostan-11-one
	Preparation B30
	(3ß,5a,25R)-3-[(heptaacetyl-ß-D-cellobiosyl)oxy]spirostan-12-one
	Preparation B31
10	(3ß,5\alpha,11\alpha,12\alpha,25R)-3-[(heptaacetyl-\beta-D-cellobiosyl)oxy]-11,12- dihydroxyspirostane
	Preparation B32
	(3ß,5ø,11ø,25R)-3-[(heptaacetyl-ß-D-cellobiosyl)oxy]-11- hydroxyspirostan-12-one
15	Preparation B33
	(3B,5a,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]spirostan-11,12-dione
	Preparation B34
20	(38,5\alpha,11\beta,12\alpha,25\beta)-3-[(heptaacetyl-\beta-D-cellobiosyl)oxy]-11,12- di(hydroxy)spirostane
	Preparation B35
	(3ß,5a,12a,25R)-3-[(ß-D-cellobiosyl)oxy]-12-hydroxyspirostan-11-one
25	Preparation B36
	(3ß,5a,12ß,25R)-3-[(heptaacetyl-ß-D-lactosyl)oxy]-12-hydroxyspirostan-11-one
	Preparation B37
	(3B,5g,25R)-3-[(dodecaacetyl-B-D-maltotriosyl)oxy]spirostan-11-one
30	Preparation B38
	(3B,5a,25R)-3-[(heptaacetyl-B-D-maltosyl)oxy]spirostan-11-one
	Preparation B39
	(1g,3ß,5g,25R)-3-[(heptaacetyl-ß-D-cellobiosyl)oxy]-1-hydroxyspirostane

WO 94/00480 PCT/US93/04092

-47-

Preparation B40

(3B,5\sigma,25R)-3-[(heptaacetyl-\textit{B-D-cellobiosyl})oxy]spirostan-6-one

5

Preparation B41

(3B,5\alpha,11B,25R)-3-[(heptaacetyl-B-D-cellobiosyl)oxy]-11-hydroxyspirostan-12-one

Preparation C1

10

15

20

25

(3B,5\sigma,25R)-3-([(heptaacetyl-B-D-lactosyl)oxy]spirostane

MERCURIC BROMIDE/MERCURIC CYANIDE PROMOTED COUPLING OF SILYLATED SPIROSTANE

Powdered 4A molecular sieves (1 g) were added to a solution of trimethylsilyl tigogenin (1.17 g, 2.4 mmol) and acetobromo lactose (3.36 g, 4.8 mmol) in CH₂Cl₂ (15 mL) and CH₃CN (5 mL) at room temperature. After stirring for 15 minutes Hg(CN)₂ (2.4 g, 9.6 mmol) and HgBr₂ (3.4 g, 9.6 mmol) were added and the mixture stirred at room temperature for three hours. The mixture was diluted with ethyl acetate (50 mL) and filtered. The filtrate was washed with 1N HCl(3 x 30 mL) and brine (1 x 30 mL), dried (Na₂SO₄) filtered and concentrated in vacuo. The product was purified by flash chromatography (10-20% EtOAc/CH₂Cl₂) to afford 400 mg product as a colorless solid. MS 489 (M+H)⁺.

¹H NMR (250 MHz, CDCl₃) δ 5.35 (d, 1H, J = 1.0 Hz); 5.2 (dd, 1H, J = 4.5, 4.5 Hz); 5.15 (dd, 1H, J = 6.0, 5.0 Hz); 4.95 (dd, 1H, J = 4.5, 1.0 Hz); 4.85 (dd, 1H, J = 5.0, 4.5 Hz); 4.55 (d, 1H, J = 6.0 Hz); 4.4 (m, 3H); 4.1 (m, 3H); 3.85 (t, 1H, J = 3.0 Hz); 3.8 (t, 1H, J = 4.5 Hz); 3.5 (m, 3H); 3.35 (t, 1H, J = 5.0 Hz); 2.15 (s, 3H); 2.12 (s, 3H); 2.07 (s, 12H); 2.0 (s, 3H); 2.0 - 0.5 (m, 27H); 0.98 (d, 3H, J = 4.0 Hz); 0.82 (s, 3H); 0.8 (d, 3H, J = 4.0 Hz); 0.73 (s, 3H).

In an analogous manner the following compounds, preparations C2-C4, were prepared from the appropriate starting material using the above general procedure.

-48-

Preparation C2

(3B,5a,25R)-3-([(heptaacetyl-B-D-maltosyl)oxy]-spirostane

5

Preparation C3

(3B,5a,25R)-3-([(triaacetyl-B-D-2-acetamido-2-deoxyglucopyranosyl)oxy]-spirostane

Preparation C4

(3B,5g,25R)-3-([(heptaacetyl-B-D-gentiobiosyl)oxy]-spirostane

10

Preparation D1

(3B,5\alpha,25R)-3-trimethylsilyloxyspirostane

SILYLATION OF SPIROSTANES

Trimethylsilyl trifluoromethanesulfonate (4 mL, 22.1 mmol) was added dropwise to a solution of tigogenin (6 g, 14.4 mmol) and triethyl amine (6 mL, 45 mmol) in CH₂Cl₂ (50 mL) at 0°C. After 1 hour, the mixture was diluted with ether (100 mL) and washed with saturated NaHCO₃ solution (2 x 50 mL) and brine (1 x 50 mL), dried (Na₂SO₄) filtered and concentrated in vacuo. Upon addition of methanol, a precipitate formed which was filtered and washed with methanol and dried to afford 6.2 g product as a colorless solid.

20

15

MP 197 -198°C. MS 489 (M + H)⁺. ¹H NMR (250 MHz, CDCl₃) δ 4.35 (q, 1H, J = 3.0 Hz); 3.5 (m, 2H); 3.4 (t, 1H, J = 5.5 Hz); 2.0-0.5 (m, 27H); 1.0 (d, 3H, J = 4.0 Hz); 0.85 (s, 3H); 0.8 (d, 3H, J = 4.0 Hz); 0.75 (s, 3H); 0.1 (s, 9H).

Preparation E1

25

$(38,5\alpha,25R)$ -3-(2-hydroxyethoxy)-spirostane

LAH REDUCTIONS

Lithium aluminum hydride (0.285 g, 7.5 mmol) was added to a solution of tigogenin-O-acetic acid ethyl ester (2.5 g, 4.98 mmol) in THF (50 mL) at 0°C. After 1 hour, the reaction was quenched by the sequential addition of H_2O (0.285 mL), 15% NaOH (0.285 mL) and H_2O (0.85 mL). The mixture was diluted with ether (25 mL) and dried with MgSO₄, filtered and concentrated in vacuo to afford 2.1 g product as a colorless solid. MP 207-208°C. MS 461 (M+H)⁺.

¹H NMR (250 MHz, CDCl₃) δ 4.4 (q, 1H, J = 3.0 Hz); 3.7 (m, 2H); 3.6 (m, 2H); 3.5 (m, 1H); 3.4 (t, 1H, J = 5.5 Hz); 3.3 (m, 1H); 2.0-0.5 (m, 28H), 1.0 (d, 3H, J = 4.0 Hz); 0.85 (s, 3H); 0.8 (d, 3H, J = 4.0 Hz); 0.75 (s, 3H).

35

WO 94/00480 PCT/US93/04092

49-

Preparation F1

((3ß,5\alpha,25R)-spirostan-3-yl)-O-acetic acid ethyl ester [Rh(OAc),], CATALYZED COUPLINGS

Ethyl diazoacetate (5.5 mL, 0.048 mol) dissolved in 30 mL of CH₂Cl₂ was added dropwise over 1 hour to a solution of tigogenin (10 g, 0.024 mol) and rhodium acetate dimer (250 mg) in CH₂Cl₂ (250 mL) at room temperature. Gas evolved throughout the addition and when the addition was complete the mixture stirred for an additional 1 hour. The mixture was diluted with hexanes (100 mL) and filtered through a plug of silica gel. The filtrate was concentrated in vacuo and upon addition of methanol to the residue, a precipitate formed which was filtered and washed with methanol and dried to afford 6.0 g product as a colorless solid. MP 119 - 120°C.MS 503 (M+H)⁺.

¹H NMR (250 MHz, CDCl₃) δ 4.35 (q, 1H, J = 3.0 Hz); 4.2 (m, 2H); 4.1 (s, 2H); 3.4 (m, 3H); 2.0 - 0.5 (m, 30H); 0.95 (d, 3H, J = 4.0 Hz); 0.8 (s, 3H); 0.75 (d, 3H, J = 4.0 Hz); 0.72 (s, 3H).

Preparation G1

(3B,5a,11B,12a,25R)-spirostan-3,11,12-triol

20

5

10

15

(3ß,5a,11a, 25R)-11,23-dibromo-3-acetoxyspirostan-12-one: The title compound was synthesized from (3ß,5a,25R)-3-acetoxyspirostan-12-one according to the procedure described in J. Chem. Soc., 1956, 4344.

(3β,5α,11α,12β,25R)-11,23-dibromospirostan-3,12-diol: (3β,5α,11α,25R)-11,23-dibromo-3-acetoxyspirostan-12-one (20.00 g, azeotropically dried with toluene) was dissolved in THF (600 mL) and cooled to -78°C. Lithium aluminum hydride (96.0 mL of 1.0 M THF solution) was slowly added and the resulting mixture was stirred at -78°C for 2 hours and 0°C for 0.5 hour. Using a cannula, the mixture was
 cautiously transferred into stirred 3 M aqueous ammonium chloride (200 mL). The organic phase was separated, combined with THF washes of the solid residues, and concentrated to give the title compound.

- (38,5a,118,128,25R)-23-bromo-11,12-epoxyspirostan-3-ol: The following procedure is a variation of that described in Helv. Act. Chim., 1953, 36, 1241.
- (3β,5α,11α,12β,25R)-11,23-dibromospirostan-3,12-diol (18.08 g) was dissolved in pyridine (500 mL) at room temperature and treated with silver oxide (70.0 g). The resulting mixture was stirred in the dark for 71 hours. The mixture was filtered and the solid washed with ether and then chloroform. These washes were combined with the filtrate and concentrated. The resulting solid was purified by flash
 chromatography (1:1 hexane:ethyl acetate) to give 12.2 g of a 1:1 mixture of the title compound and (3β,5α,25R)-23-bromospirostan-3-ol-12-one. Further chromatography (7:3 hexane:ethyl acetate) provides pure title compound.
 - (38,5a,118,12a,25R)-23-bromo-12-(trichloroacetoxy)spirostan-3,11-diol: Using the procedure described in <u>J. Chem. Soc.</u>, 1956, 4330, (38,5a,118,128,25R)-23-bromo-11,12-epoxyspirostan-3-ol was treated with trichloroacetic acid in toluene at room temperature for 3 days to give the title compound.
 - (3ß,5α,11β,12α,25R)-23-bromo-spirostan-3,11,12-triol: Using the procedure
 described in <u>J. Chem. Soc.</u>, 1956, 4330, (3ß,5a,11ß,12α,25R)-23-bromo-12(trichloroacetoxy)spirostan-3,11-diol was saponified with sodium hydroxide in water and ethanol to give the title compound.
 - (3β,5α,11β,12α,25R)-spirostan-3,11,12-triol: Using the procedure described in J.
 Chem. Soc., 1956, 4330, (3β,5a,11β,12α,25R)-23-bromo-12-(trichloroacetoxy)-spirostan-3,11-diol was reduced with zinc and acetic acid to give the title compound.

Preparation G2 (3B,5a,12a,25R)spirostan-3,12-diol-11-one

30

(3β,5α,11β,12α,25R)-3,12-di(acetoxy)spirostan-11-ol: Using the procedure described in <u>J. Chem. Soc.</u>, 1956, 4330, (3β,5α,11β,12α,25R)-spirostan-3,11,12-triol (preparation G1) was selectively acetylated with acetic anhydride and pyridine to give the title compound.

WO 94/00480 PCT/US93/04092

-51-

(3β,5α,12α,25R)-3,12-di(acetoxy)spirostan-11-one: Using the procedure described in Org. Syn., 1976, 55, 84, (3β,5α,11β,12α,25R)-3,12-di(acetoxy)-spirostan-11-ol was oxidized with chromium trioxide and pyridine in methylene chloride to give the title compound.

(38,5a,12a,25R)-spirostan-3,12-diol-11-one: Using the procedure described in Syn., 1973, 790, (36,5a,12a,25R)-3,12-di(acetoxy)spirostan-11-one was saponified with potassium cyanide in water, methanol and THF to give the title compound.

<u>Preparation G3</u> (38,5\sigma,116,25R)spirostan-3,11-diol

15 (38,5a,118,25R)spirostan-3,11-diol: (38,5a,25R)spirostan-3-ol-11-one (Aldrich Chemical Company, Milwaukee, WI or Steraloids Inc., Wilton, N.H., or see preparation G13) was converted into the title compound via reduction with lithium aluminum hydride in THF at room temperature according to the procedure described in J. Am. Chem. Soc., 1951, 73, 1777.

20

25

Preparation G4 (36,5\sigma,11\sigma,25R)spirostan-3,11-diol

(3ß,5a,11a,25R)spirostan-3,11-diol: (3ß,5a,25R)spirostan-3-ol-11-one (Aldrich Chemical Company, Milwaukee, WI or Steraloids Inc., Wilton, N.H., or see preparation G13) was converted into the title compound via reduction with lithium and ammonia according to the procedure described in J. Am. Chem. Soc., 1953, 75, 1282.

Preparation G5

30

35

1777.

(3B,5a,11B,12B,25R)spirostan-3,11,12-triol

(3β,5α,11β,12β,25R)spirostan-3,11,12-triol: (3β,5α,12β,25R)-3,12-di(acetoxy)spirostan-11-one (purchased from Steraloids, Inc., or see preparation G13) was converted into the title compound via reduction with lithium aluminum hydride in THF at room temperature according to the procedure described in J. Am. Chem. Soc., 1951, 73,

Preparation G6

(3B.5a.11a.12B.25R)spirostan-3.11.12-triol

5

15

- (3ß,5\alpha,12ß,25R)spirostan-3,12-diol-11-one: (3ß,5\alpha,12ß,25R)-3,12-di(acetoxy)spirostan-11-one (purchased from Steraloids, Inc., or see preparation G13) was saponified with potassium carbonate in water, methanol and THF to provide the title compound.
- 10 (3β,5α,11α,12β,25R)spirostan-3,11,12-triol: (3β,5α,12β,25R)spirostan-3,12-diol-11-one was converted into the title compound via reduction with lithium and ammonia according to the procedure described in J. Am. Chem. Soc., 1953, 75, 1282.

Preparation G7

(3B,5a,12a,25R)spirostan-3,12-diol

(3β,5α,12α,25R)spirostan-3,12-diol: Using the procedure described in <u>J. Am. Chem. Soc.</u>, 1954, <u>76</u>, 4013, (3β,5α,25R)-spirostan-3-ol-12-one was reduced with lithium aluminum hydride in ether to give a mixture of C-12 alcohols from which the title compound was isolated.

Preparation G8 (36,5\alpha,25R)-spirostan-3-ol-11,12-dione

- 25 (3β,5α,12β,25R)-3-(t-butyldimethylsilyloxy)spirostan-12-ol-11-one: Using the procedure described in J. Am. Chem. Soc., 1972, 94, 6190, (3β,5α,12β,25R)-spirostan-3,12-diol-11-one (see preparation G6) was silylated with t-butyldimethylchlorosilane and imidazole in DMF to give the title compound.
- 30 (38,5a,25R)-3-(t-butyldimethylsilyloxy)spirostan-11,12-dione: Using the procedure described in Org. Syn., 1976, 55, 84, (38,5a,128,25R)-3-(t-butyldimethylsilyloxy)-spirostan-12-ol-11-one was oxidized with chromium trioxide and pyridine in methylene chloride to give the title compound.

(3β,5σ,25R)-spirostan-3-ol-11,12-dione: Using the procedure described in <u>J. Am. Chem. Soc.</u>, 1972, 94, 6190, (3β,5a,25R)-3-(t-butyldimethylsilyloxy)spirostan-11,12-dione was desilylated with hydrofluoric acid in acetonitrile to give the title compound.

Preparation G9 (38.5 σ ,118,25R)-spirostan-3,11-diol-12-one

(3β,5α,11β,12β,25R)-3-(t-butyldimethylsilyloxy)spirostan-11,12-diol: (3β,5α,12β,25R)-3-(t-butyldimethylsilyloxy)spirostan-12-ol-11-one (see procedure G8) was converted into the title compound via reduction with lithium aluminum hydride in THF at room temperature according to the procedure described in J. Am. Chem. Soc., 1951, 73, 1777.

15

(38,5\alpha,118,128,25R)-3-(t-butyldimethylsilyloxy)-12-acetoxyspirostan-11-ol: (38,5\alpha,118,128,25R)-3-(t-butyldimethylsilyloxy)spirostan-11,12-diol was selectively acetylated with acetic anhydride, pyridine and dimethylaminopyridine in methylene chloride to give the title compound.

20

(38,5\alpha,118,128,25R)-3-(t-butyldimethylsilyloxy)-11-(trimethylsilyloxy)-12-acetoxyspirostane: (38,5\alpha,118,128,25R)-3-(t-butyldimethylsilyloxy)-12-acetoxyspirostan-11-ol was silylated with trimethylsilyltriflate and 2,6-lutidine in methylene chloride according to the procedure described in Tetrahedron Letters, 1981,

25 <u>22,</u> 3455.

(38,5\u03ba,118,128,25R)-3-(t-butyldimethylsilyloxy)-11-(trimethylsilyloxy)spirostan-12-ol: (38,5\u03ba,118,128,25R)-3-(t-butyldimethylsilyloxy)-11-(trimethylsilyloxy)-12-acetoxyspirostane was deacetylated by treatment with lithium aluminum hydride in THF followed by catious addition aqueous ammonium chloride. The resulting title compound suffered 11 to 12 silyl group migration on silica gel, and thus had to used unpurified.

20

25

30

35

(38,5a,118,25R)-3-(t-butyldimethylsilyloxy)-11-(trimethylsilyloxy)spirostan-12-one: (38,5a,118,128,25R)-3-(t-butyldimethylsilyloxy)-11-(trimethylsilyloxy)spirostan-12-ol was oxidized with chromium trioxide and pyridine in methylene chloride according to the procedure described in Org. Syn., 1976, 55, 84 to give the title compound.

 $(3\beta,5\alpha,11\beta,25R)$ -spirostan-3,11-diol-12-one: The title compound was synthesized from $(3\beta,5\alpha,11\beta,25R)$ -3-(t-butyldimethylsilyloxy)-11-(trimethylsilyloxy)spirostan-12-one was desilylated with hydrofluoric acid in acetonitrile according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1972, <u>94</u>, 6190. The title compound must be carefully handled because it will rearrange to $(3\beta,5\alpha,12\beta,25R)$ -spirostan-3,12-diol-11-one if exposed to base.

Preparation G10

(3B,5g,11g,25R)spirostan-3,11-diol-12-one

(3ß,5\u03b3,11\u03b3,25R)3,11-di(acetoxy)spirostan-12-ol: (3ß,5\u03b3,11\u03b3,11\u03b3,25R)-spirostan-3,11,12-triol (see preparation G6) was acetylated according to the procedure described in <u>J. Am. Chem. Soc.</u>, 1955, <u>77</u>, 1632 to give a mixture of acetates from which the title compound could be isolated.

(3β,5α,11α,25R)3,11-di(acetoxy)spirostan-12-one: (3β,5α,11α,12β,25R)3,11-di(acetoxy)spirostan-12-ol was oxidized with chromium trioxide and pyridine in methylene chloride according to the procedure described in Org. Syn., 1976, 55, 84 to give the title compound.

(3B,5\alpha,11\alpha,25R)spirostan-3,11-diol-12-one: (3B,5\alpha,11\alpha,25R)-3,:1-di(acetoxy)spirostan-12-one was saponified with sodium methoxide in methanol and THF to give the title compound.

Preparation G11

(3B,5a,11a,12a,25R)spirostan-3,11,12-triol

(3B,5a,25R)spirostan-3-ol-12-tosylhydrazone: (3B,5a,25R)-spirostan-3-ol-12-one (8.00g) was dissolved in glacial acetic acid (200 mL) and warmed to 50°C. Paratoluenesulfonylhydrazide (6.928 g) was added and the solution was stirred at 50°C for

30 min. After an additional 2 hours of stirring at room temperature, water (200 mL) was added. The resulting solid was collected, washed with water (100 mL), dried, triturated 5 with refluxing acetone (300 mL), filtered hot and dried to give 3.903 g of the title compound.

 $(3\beta,5\alpha,25R)$ spirost-11-en-3-ol: A mixture of $(3\beta,5\alpha,25R)$ spirostan-3-ol-12-tosylhydrazone (9.100 g) and sodium methoxide (8.379 g) in DMF (200 mL) was heated to 150°C for 10 35 minutes, then cooled to room temperature. The mixture was then poured into ice water (1200 mL) and the resulting suspension filtered. The collected solid was washed with water (100 mL), air-dried, and dissolved in methylene chloride (700 mL). This solution was washed with water (2 X 200 mL), dried with MgSO4, and concentrated to give a white solid. Following flash chromatography, 2.384 g of the title compound (mp 179-181 °C, lit. 188-192 °C - <u>J. Am. Chem. Soc.</u>, 1954, <u>76,</u> 4013) was isolated.

(38,5a,11a,12a,25R)spirostan-3,11,12-triol: (38,5a,25R)spirost-11-en-3-ol was oxidized to the title compound with osmium tetroxide and N-methylmorpholine-N-oxide in water, t-butanol and acetone according to the procedure describe in Tetrahedron Letters, 1976, 1973.

Preparation G12

(3B,5a,12B,25R)spirostan-3,12-diol-11-one

(3B,5a,11B,25R)-11-bromospirostan-3-ol-12-one: A glass lined reactor was charged with 50 gallons of methanol then subsurface sparged with hydrochloric acid gas until 7.7 Kg (5.0 eq) were charged. Upon completion of this sparge, the reactor was charged with 18.8 Kg (42.2 mole) of (38,5a,25R)spirostan-3-ol-12-one, 50 gallons of methanol and 10 gallons of methylene chloride. This mixture was cooled to 10 °C and a solution of 8.4 Kg bromine (52.7 mole, 1.25 eq) in 10 gallons of methylene chloride was added over 2 hours while a pot temperature of approximately 10°C was maintained. Once the addition was complete the reaction was allowed to warm to room temperature and was stirred for 2 hours. TLC at this point indicated complete reaction.

The reaction was diluted with 50 gallons of water and stirred for 10 minutes. After separation of layers, the aqueous layer was extracted twice with 30 gallons of methylene chloride. The three combined organic extracts were washed twice with 30

15

20

25

gallons of water, once with 30 gallons of saturated brine, then dried using 7.0 Kg of magnesium sulfate. The drying agent was removed by filtration on a 30 inch Lapp followed by two 3 gallon methylene chloride washes. The filtrate and washes combined were atmospherically distilled to a 7 gallon total volume. Two 10 gallon methanol charges were made followed by continued distillation. When a final volume of <10 gallons had been reached the mixture was cooled to room temperature. The resulting suspension was granulated for 2 hours, filtered on a 30 inch Lapp, and the filter cake was washed twice with 3 gallons of methanol. Vacuum drying the filter cake at 45-50° C yielded 12.6 Kg (58.6% yield) of the title compound.

(38,5a,128,25R)spirostan-3,12-diol-11-one: A glass lined reactor was charged with 12.4 Kg of (38,5a,118,25R)-11-bromospirostan-3-ol-12-one (24.34 mole), 33 gallons of t-butanol, 33 gallons of water and 7.5 Kg (189 mole, 7.75 eq) of sodium hydroxide pellets. The reaction was heated to reflux over 1.5 hours, maintained at reflux for 4.5 hours (pot temperature was 83° C), then cooled to room temperature. TLC at this point indicated complete reaction.

The reaction was distilled to remove the t-butanol. This was accomplished both by vacuum and atmospheric distillation. During the concentration two 32.5 gallon charges of water were added. Once the t-butanol had been removed, the aqueous suspension was cooled to room temperature and granulated for 2 hours. The suspension was filtered on a 30 inch Lapp, washed twice with 3 gallons of water, and the filter cake was air dried at 60 °C. This afforded 11.1 Kg of the title compound.

25

30

35

20

10

15

Preparation G13 (38,5\sigma,25R)spirostan-3-ol-11-one

(3β,5α,12β,25R)-3,12-diacetoxyspirostan-11-one: Aglass lined reactor was charged with 26 gallons of pyridine, 26 gallons of acetic anhydride and 11.0 Kg of (3β,5α,12β,25R)spirostan-3,12-diol-11-one (preparation G12). This mixture was refluxed for 2 hours (pot temperature 128° C) and allowed to cool to room temperature. The reaction was vacuum distilled to a total volume of 15 gallons (pot temperature approximately 45° C during distillation). The suspension was diluted with 25 gallons of acetic acid and further vacuum distilled to a 15 gallon total volume (pot temperature

WO 94/00480 PCT/US93/04092

at end approximately 80° C). The mixture was diluted with 87 gallons of water and cooled to room temperature. After 5 hours of granulation, the titled compound was 5 isolated by filtration on a 30 inch Lapp followed by two 3 gallon water washes. The filter cake was dried at 60° C under vacuum to yield 12.2 Kg (93.3%).

(3B,5\sigma,25R)spirostan-3-ol-11-one: A stainless steel reactor was cooled to -80° C by passing liquid nitrogen through internal coils. Ammonia was added to the reactor until 54.5 Kg (80 liters, 3,200 mole, 170 eq) had been charged.

At the same time that the ammonia charge was commencing, a glass lined reactor was charged with 10.0 Kg of (3B,5a,12B,25R)-3,12-diacetoxyspirostan-11-one 18.84 mole) and 40 galions of THF. This solution was atmospherically distilled until a 26 gallon total volume had been reached.

At the completion of the ammonia charge, 2.8 Kg of calcium turnings (69.0 gram atoms, 3.7 eq) were added over 30 minutes while maintaining a pot temperature of -50° C. At the completion of this addition the THF solution of (3B,5\alpha,12B,25R)-3,12diacetoxyspirostan-11-one was added over 20 minutes (pot temperature at the end of the addition was -35° C) followed by a 1.0 gallon THF rinse. The reaction mixture was stirred for 30 minutes at - 35°C to -40°C. While the reaction was at -35° C to -40° C, 3.33 liters of bromobenzene (4.98 Kg, 31.7 mole, 1.68 eq) were added followed by 3.33 liters of water.

After this addition, the distillation of ammonia from the reactor was initiated. This distillation was directed to a water scrubber. Once all of the ammonia had been 25 removed, the reaction (now at 24° C) was transferred to a glass lined reactor followed by a 4 gallon THF rinse. The solution and rinse combined were vacuum distilled to a thick oil. To this was added 35 gallons of methanol and 3.3 Kg (59 mole) of potassium hydroxide pellets. This mixture was heated at reflux for 1 hour, cooled, then 10 liters of acetic acid and 44 gallons of water were charged. This suspension was further cooled to room temperature and granulated for 1 hour. The titled compound was isolated by filtration on a 30 inch Lapp followed by a 5 gallon 3:1 water/methanol wash. Vacuum drying at 55° C yielded 7.05 Kg (86.9%).

30

10

15

Preparation G: Physical Data

Satisfactory MS and IR data was obtained on all of the (3ß,5\alpha,25R)spirostan-3ols described in preparation G (see table 1). The various diol and triol products could be distinguished by proton NMR (see table 2)

Table 1: Diagnostic Mass Spectrometry and Infrared Data

10	compound	molecular formula	LSMIS parent ion (m/z)	IR diagnostic resonances (cm-1, intensity, solvent)
10	44 al	C 11 0	422	2575 (m) 2440 (m) (CHCL)
	11 <i>a-</i> ol	C ₂₇ H ₄₄ O ₄	433	3575 (m), 3440 (m) (CHCl ₃)
	11B-oi	C27H44O4	433	3560 (m), 3425 (m) (CHCl ₃)
	12 <i>a</i> -ol	C27H44O4	433	3590 (m), 3420 (m) (CHCl ₃)
	12B-ol	C27H44O4	433	
	11 <i>a</i> ,12 <i>a</i> -diol	C ₂₇ H ₄₄ O ₅	449	3424 (m) (KBr)
	11a,12B-diol	C27H44O5	449	3550 (m), 3450 (m) (CHCl ₃)
15	11β,12 <i>σ</i> -diol	C ₂₇ H ₄₄ O ₅	449	3441 (m) (KBr)
	11B,12B-diol	C ₂₇ H ₄₄ O ₅	449	3600 (m), 3450 (m) (CHCl ₃)
	11 <i>α</i> -ol-12-one	C ₂₇ H ₄₂ O ₅	447	3515 (m), 1705 (s) (KBr)
	11ß-ol-12-one	C ₂₇ H ₄₂ O ₅	447	3450 (m), 1712 (s) (KBr)
	12 <i>α</i> -ol-11-one	C ₂₇ H ₄₂ O ₅	447	3410 (m), 1706 (s) (KBr)
	12ß-ol-11-one	C ₂₇ H ₄₂ O ₅	447	3475 (m), 1708 (s) (CHCL)
20	11,12-dione	C ₂₇ H ₄₀ O ₅	445	3600 (w), 3400 (m), 1710 (w) 1670 (s), 1605 (m) (CHCl ₃) ¹
	11-one	C ₂₇ H ₄₂ O ₄	431	3600 (w), 3450 (m), 1705 (s) (CHCL)

^{1 -} IR data suggest that this compound readily tautomerizes enol ketone form in CHCl₃.

25 Table 2: Diagnostic Proton Nuclear Magnetic Resonance Data²

	compound	peaks >2 ppm
	11 <i>a</i> -ol	3.90 (ddd, 6,6 &4 Hz, 1H), 2.26 (dt, 13&4, 1H)
	11B-ol	4.22 (br s, 1 H)
	12 <i>a</i> -ol	3.67 (s, 1 H), 2.37 (dd, 8 & 7 Hz, 1H)
	12B-ol	3.26 (dd, 10 & 4 Hz, 1H)
30		
	11 <i>a</i> ,12 <i>a</i> -diol	3.91(m, 1H), 3.56 (d, 3H, 1H), 2.45 (dd, 9 &7 Hz, 1H)
	11a,12G-diol	3.55(m, 1H), 3.03 (d, 8H, 1H), 2.21 (dt, 12 &4 Hz, 1H)
	11B,12a-diol	4.12 (br s, 1H), 3.55 (d, 2 Hz, 1H), 2.36 (dd, 9 &7 Hz, 1H)
	11B,12B-diol	4.07 (br s, 1H), 3.13 (d, 3 Hz, 1H)
	11 <i>a</i> -ol-12-one	3.72 (m, 1H), 2.39 (dt, 13 & 4 Hz, 1H)
	11B-ol-12-one	3.96 (m, 1H), 2.2 (m, 1H)
35	12 <i>α</i> -ol-11-one	3.51 (s, 1H), 2.57 (dd, 8 & 7 Hz, 1H), 2.2 (complex, 7H)
00	12ß-ol-11-one	3.78 (s, 1H), 2.39 (dt,13 &4 Hz, 1H), 2.1 (m, 2H)

2 - All samples run in CDCl₃ except 11 θ -ol-12-one which was run in DMSO-d_e. Peaks for H_{1e}, H₃, H_{26 ∞} and H_{26 ∞} are also observed at >2 ppm. In CDCl₃, these are observed at 4.37 (ddd, J = 9, 9 & 7 Hz, 1H), 3.56 (heptet, J = 4 Hz, 1H), 3.45 (ddd, J = 10, 6 & 2 Hz, 1H), 3.35 (t, J = 11 Hz, 1H).

Preparation H1

(5a,25R)-spirostan-3-one

Pyridinium chlorochromate (PCC) was added to a mixture of tigogenin (50.00 g, 120.0 mmol), celite (160 g), in CH₂Cl₂ (1000 mL) at 0° C. The reaction was allowed to come to ambient temperature and was stirred for 5 hours. The reaction was diluted with 1000 4 mL Et₂O and was filtered through a silica gel plug. The plug was washed with an additional 6000 mL Et₂O. The filtrate was concentrated in vacuo to afford 45.00 g of the title compound (90.4%).

¹HNMR (250 MHz, CDCl₃) d 4.38 (q, J = 7Hz, 1H); 3.40 (m, 2H); 2.20-2.45 (m, 3H); 0.70-2.14 (m, 36H); 1.02 (s, 3H); 0.96 (d, J = 7 Hz, 3H); 0.76 (s, 3H); 0.76 (d, J = 7Hz, 3H). MS: 415 (M + H) $^+$; MP 209-211 $^{\circ}$ C.

Preparation H2

(2a,5a,25R)-2-bromospirostan-3-one

A mixture of (5a,25R)-spirostan-3-one (1.00 g, 2.41 mmol) and tetrahydrofuran (10 mL) was cooled to -78°C under nitrogen atmosphere. Bromine was added (0.39 g, 2.41 mmol) and the reaction mixture was gradually warmed to room temperature. After three hours, the reaction was quenched by the addition of saturated sodium bisulfite solution. The mixture was diluted with ethyl acetate, washed with saturated sodium bisulfite solution (1x), saturated sodium bicarbonate (1x), brine (1x), dried (sodium sulfate), filtered, and concentrated in vacuo. Upon addition of ether, a precipitate formed which was filtered and washed with hexane: to give 1.20 g, (85%) of the title compound.

¹HNMR (250 MHz, CDCl₃) d 4.75 (q, J = 7Hz, 1H); 4.40 (q, J = 7Hz, 1H); 3.40 (m, 2H); 2.64 (q, J = 6, 1H); 2.40 (m, 2H); 0.70-2.55 (m, 34H); 1.10 (s, 3H); 0.96 (d, J = 7Hz, 3H); 0.80 (s, 3H); 0.80 (d, J = 7, 3H). MS 493 (M + H)⁺.

15

20

25

10

15

20

25

30

35

Preparation H3

(5a,25R)-spirost-1-en-3-one

A mixture of lithium bromide (0.700 g, 8.06 mmol), lithium carbonate (1.20 g, 16.24 mmol) and anhydrous N,N-dimethylformamide (30 mL) were heated under nitrogen atmosphere to 95°C. To this mixture, (2\alpha,5\alpha,25R)-2-bromospirostan-3-one (4.00 g, 8.11 mmol) was added. The reaction mixture was stirred at 130°C for 3 hours. After cooling to room temperature, the reaction mixture was diluted with ethyl acetate, washed with water (3x), brine (1x), dried (sodium sulfate), filtered and concentrated in vacuo to afford 3.31 g, (98%) of the title compound.

¹HNMR (250 MHz, CDCl₃) d 7.10 (d, J = 10Hz, 1H); 5.85 (d, J = 10Hz, 1H), 4.40 (q, J = 7Hz, 1H); 3.40 (m, 2H); 2.30 (m, 2H); 0.70-2.05 (m, 33H); 1.02 (s, 3H); 0.96 (d, J = 7Hz, 3H); 0.80 (s, 3H); 0.78 (d, J = 7Hz, 3H). MS 413 (M + H) $^+$.

Preparation H4

$(1\alpha, 2\alpha, 5\alpha, 25R)$ -1,2-epoxy-spirostan-3-one

A mixture of (5a,25R)-spirost-1-en-3-one (2.87 g, 6.96 mmol), tetrahydrofuran (30 mL), methanol (50 mL) and 15% sodium hydroxide (1 mL) was stirred under nitrogen atmosphere. The mixture was cooled to 0°C and 30% hydrogen peroxide (5 mL) was added. The reaction mixture was gradually warmed to room temperature and stirred for 4 hours. The reaction was diluted with ethyl acetate, cooled to 0°C, and then quenched with saturated sodium bisulfite solution. The mixture was washed with saturated sodium bisulfite (2x), brine (1x), dried (sodium sulfate), filtered and concentrated in vacuo to give 2.64 g, (88%) of the title compound.

¹HNMR (250 MHz, CDCl₃) d 4.40 (q, J = 7Hz, 1H); 3.40 (m, 3H); 3.24 (d, J = 6Hz, 1H); 2.25 (dd, J = 18, 4 Hz, 1H); 0.70-2.28 (m, 34H); 0.98 (d, J = 7Hz, 3H); 0.92 (s, 3H); 0.80 (s, 3H); 0.78 (d, J = 7Hz, 3H). MS 429 (M + H)⁺.

Preparation H5

$(1\alpha,3\beta,5\alpha,25R)-1,3-di(hydroxy)$ spirostane

Lithium aluminum hydride (0.43 g, 15.38 mmol) was added to a solution of (1\alpha,2\alpha,5\alpha,25R)-1,2-epoxy-spirostan-3-one in THF (20 mL) at 0°C. The reaction was gradually warmed to room temperature and after 3 hours, additional lithium aluminum hydride (0.10 g, 3.58 mmol) was added. After 1 hour, the reaction was

WO 94/00480 PCT/US93/04092

-61-

cooled to 0°C and quenched by the sequential addition of H₂O (0.75 mL), 15% NaOH (0.75 mL), and H₂O (1.50 mL). The mixture was dried with MgSO₄, filtered, and concentrated in vacuo. The product was purified by flash chromatography (50% EtOAc/50% hexane to 95% EtOAc/5%MeOH) to afford 0.460 g, (34%) of the title compound.

¹HNMR (250 MHz, CDCl₃) d 4.48 (q, J = 7Hz, 1H); 4.04 (m, 1H); 3.80 (m, 1H); 3.40 (m, 2H); 0.75-2.05 (m, 37H); 0.96 (d, J = 6Hz, 3H); 0.84 (s, 3H); 0.78 (d, J = 6Hz, 3H); 0.76 (s, 3H). MS 433 (M + H)⁺.

Preparation I1

(3β,5σ,25R)-3-[(heptaacetyl-β-D-cellobiosyl)oxy]spirostan-1-one

Pyridinium chlorochromate (0.123, 0.57 mmol) was added to a mixture of (1α,3β,5α,25R)-3-[(heptaacetyl-β-D-cellobiosyl)oxy]-1-hydroxyspirostane (0.2 g, 0.19 mmol, see preparation B39), celite (0.2 g), in CH₂Cl₂ (5 mL) at 0°C. The reaction was allowed to come to ambient temperature and was stirred for 2 hours. The reaction was diluted with 15 mL Et₂O and was filtered through a silica gel plug. The plug was washed with an additional 500 mL Et₂O. The filtrate was concentrated in vacuo to afford 0.18 g of the title compound (90%).

¹HNMR (250 MHz, CDCl₃) d 5.15 (m, 3H); 4.90 (m, 2H); 4.50 (m, 2H); 4.35 (m, 2H); 4.05 (m, 2H); 3.65 (m, 3H); 3.40 (m, 2H); 2.35 (t, J = 12.5 Hz, 1H); 2.60 (q, J = 6 Hz, 1H); 1.95-2.20 (m, 21H); 0.70-1.90 (m, 37H); 1.15 (s, 3H); 0.95 (d, J = 7 Hz, 3H); 0.80 (d, J = 6 Hz, 3H); 0.76 (s, 3H). MS: 1049 (M + H)⁺.

25

30

10

15

20

Preparation J1

(3ß,25R)-3-ethoxymethoxy-5-spirostene

A mixture of diosgenin (2.5 g, 6.0 mmol), chloromethyl ethyl ether (1.14 g, 12.0 mmol), diisopropylethylamine (3.90 g, 30.0 mmol) and 1,2-dichloroethane (75 mL) was stirred under nitrogen atmosphere at ambient temperature for 4 hours. Methanol (<1 mL) was added to quench the reaction. The mixture was diluted with ethyl acetate and washed with water (2x), brine (1x), dried over sodium sulfate, filtered and concentrated in vacuo to give 2.17 g (76.5%) of the title compound as a colorless solid.

20

25

30

35

¹H NMR (250 MHz, CDCl₃) d 5.35 (d, 2H, J=7.0 Hz); 4.75 (s, 2H); 4.4 (m, 1H);3.6 (q, 2H, J=7.0 Hz); 3.4 (m, 2H); 3.35 (t, 1H, J=11 Hz); 2.4-0.7 (m, 38H); 1.2 (s, 3H); 0.95 (d, 3H, J=7 Hz); 0.8 (d, 3H, J=7 Hz); 0.75 (s, 3H). MS: 777 (M+Na)+; mp 125-127°C.

Preparation J2

(3B,5a,6a,25R)-3-ethoxy methoxy-6-hydroxyspirostane

Borane-tetrahydrofurancomplex (0.68 mL, 0.68 mmol) was added to a solution of (3B,25R)-3-ethoxymethoxy-5-spirostene (0.10 g, 0.21 mmol) in tetrahydrofuran (8 10 mL). The mixture was stirred under nitrogen atmosphere at ambient temperature for 3.5 hours. The reaction mixture was cooled to 0°C and methanol (1.5 mL), 15% sodium hydroxide solution (1.5 mL) and 30% hydrogen peroxide (1.5 mL) were added. The reaction mixture was then gradually warmed to ambient temperature and stirred overnight. The reaction was quenched at 0°C by the addition of saturated sodium bisulfite solution. The mixture was diluted with ethyl acetate and washed with ammonium chloride solution (1x), brine (1x), dried (sodium sulfate), filtered and concentrated in vacuo to give 0.11 g of a mixture of the 6a-alcohol and the 6Balcohol. The two products were separated by flash chromatography on silica gel (6:4 hexane/ethyl acetate). The major product (Rf=0.40) was identified to be the title compound.

> ¹H NMR (250 MHz; CDCl₃) d 4.7 (s, 2H); 4.4 (m, 1H); 3.6 (q, 2H, J=11.0 Hz);3.45 (m, 3H); 3.35 (t, 1H, J=11.0 Hz); 2.3-0.6 (m, 41H); 1.8 (d, 3H, J=7.0 Hz);0.82 (s, 3H), 0.78 (d, 3H, J=7.0 Hz); 0.75 (s, 3H). MS: 491 (M+H)⁺; mp 171°C.

Preparation J3

(3B,5a,25R)-3-ethoxymethoxy-spirostan-6-one

Pyridinium chlorochromate (1.98 g, 9.20 mmol) was added to a mixture of (3B,5a,6a,25R)-3-ethoxy methoxy-6-hydroxyspirostane (0.90 g, 1.8 mmol) and celite (8.0 g) in anhydrous dichloromethane at 0°C. The reaction mixture was gradually warmed to ambient temperature over 1 hour and allowed to stir for an additional 5 hours. The reaction mixture was then filtered through a plug of silica gel using ether as the eluent. The combined ether fractions were concentrated in vacuo to afford 0.80 g (91%) of the title compound as a colorless solid.

20

25

30

35

¹H NMR (250 MHz; CDCl₃) d 4.75 (m, 2H); 4.4 (m, 1H); 3.6 (q, 2H, J=7.0 Hz); 3.45 (m, 2H); 3.35 (t, 1H, J=11.0 Hz); 2.4-0.6 (m, 40H); 0.9 (d, 3H, J=7.0 Hz); 0.8 (d, 3H, J=7.0 Hz); 0.78 (s, 6H). MS: 489.0 (M+H)⁺; mp 191-193°C.

Preparation J4

(3B.5a.25R)-spirostan-6-one

Concentrated hydrochloric acid (2 drops) was added to a solution of

(36,5a,25R)-3-ethoxymethoxy-spirostan-6-one (0.70 g, 1.43 mmol) in methanol (10 mL) and tetrahydrofuran (10 mL). The mixture was stirred under nitrogen atmosphere and heated to 62°C. After 15 minutes, the reaction was cooled to 0°C and neutralized with 15% sodium hydroxide solution. The mixture was concentrated in vacuo then diluted with ethyl acetate. The organic layer was washed with water

(2x), brine (1x), dried (sodium sulfate), concentrated in vacuo and purified by flash chromatography (1:1 hexane/ethyl acetate) to afford 0.55 g (89.4%) of the title compound as a colorless solid.

¹H NMR (250 MHz; CDCl₃) d 4.4 (m, 1H); 3.45 (m, 2H); 3.35 (t, 1H,J=11.0 Hz); 2.35-0.6 (m, 38H); 0.95 (d, 3H, J=7.0 Hz); 0.75 (d, 3H, J=7.0); 0.71 (s, 6H). MS: 431 (M+H)⁺; mp 210-212°C.

Preparation K1

(36,5\(\alpha\),6\(\alpha\),25R)-3-[(heptaacetyl-\(\beta\)-D-cellobiosyl)oxy] 6-hydroxyspirostane Sodium borohydride (0.11 g, 2.86 mmol) was added to a solution of (3\(\beta\),5\(\alpha\),25R)-3-[(heptaacetyl-\(\beta\)-D-cellobiosyl)oxy] spirostan-6-one (1.5 g, 1.43 mmol, see preparation B41) in ethanol (20 mL) and dichloromethane (5 mL) and the mixture was stirred under nitrogen atmosphere at room temperature for 4 hours. The reaction mixture was then cooled to 0°C and was neutralized with 1N hydrochloric acid. The mixture was partially concentrated in vacuo and then was diluted with ethyl acetate, washed with 1N hydrochloric acid (1x), brine (1x), dried (sodium sulfate), filtered and concentrated in vacuo to afford 1.00 g (66%) of the title compound as a colorless solid.

¹H NMR (250 MHz; CDCl₃) d 5.2-4.4 (m, 14H); 4.3 (m, 1H); 3.75-3.35 (m, 3H); 3.3 (t, 1H, J=11.0 Hz); 2.15-0.5 (m, 59H); 0.95 (s, 3H); 0.90 (d, 3H, J=7.0 Hz); 0.75 (s, 3H); 0.70 (d, 3H, J=7.0 Hz). MS: 1051 (M+H)⁺

It should be understood that the invention is not limited to the particular embodiments shown and described herein, but that various changes and modifications may be made without departing from the spirit and scope of this novel concept as defined by the following claims.

10

15

20

25

Claims

5 1. A spirostanyl glycoside of Formula IA

_

Q⁴ and Q⁵ are both methylene;

and wherein

R1 is

B-D-glucopyranosyl,

B-D-glucopyranuronosyl,

B-D-2-acetamido-2-deoxy-glucopyranosyl,

35
B-D-galactopyranosyl,

```
B-D-fucopyranosyl,
          B-L-fucopyranosyl,
          B-D-xylopyranosyl,
5
          B-L-xylopyranosyl,
          q-D-arabanopyranosyl,
          a-L-arabanopyranosyl,
          a-D-cellobiosyl,
          B-D-cellobiosyl,
10
          B-D-lactosyl,
          B-D-maltosyl,
          B-D-gentiobiosyl,
          3-O-B-D-galactopyranosyl-a-D-arabanopyranosyl or
          B-D-maltotriosyl;
15
                                            or (B):
```

 Q^1 , Q^4 and Q^5 are all methylene;

$$H$$
 OR^1 R^1O H R^1O -alkylene(C_2 - C_3)- O H Q^3 is $-C$ -, $-C$ -

25

or

30 and wherein

R1 is

B-D-glucopyranosyl,

B-D-glucopyranuronosyl,

B-D-2-acetamido-2-deoxy-glucopyranosyl,

35 β-D-fucopyranosyl,

B-D-cellobiosyl,

```
B-L-fucopyranosyl,
           B-D-xylopyranosyl,
           B-L-xylopyranosyl,
 5
           a-D-arabanopyranosyl,
           a-L-arabanopyranosyl,
           B-D-cellobiosyl,
           B-D-lactosyl,
           B-D-maltosyl,
10
           B-D-gentiobiosyl,
           3-O-B-D-galactopyranosyl-a-D-arabanopyranosyl or
           B-D-maltotriosyl;
                                                 or (C):
15
           Q1, Q4 and Q5 are all methylene;
           Q<sup>2</sup> is carbonyl;
                                          R<sup>1</sup>0-alkylene(C<sub>2</sub>-C<sub>3</sub>)-0 H
20
     OF
            R<sup>1</sup>0-alkylene(C<sub>2</sub>-C<sub>3</sub>)-0
           C<sub>25</sub> is (R);
25
      and wherein
           R1 is
           B-D-glucopyranuronosyl,
           B-D-2-acetamido-2-deoxy-glucopyranosyl,
           B-D-fucopyranosyl,
30
           B-L-fucopyranosyl,
           B-D-xylopyranosyl,
           B-L-xylopyranosyl,
           a-D-arabanopyranosyl,
           a-L-arabanopyranosyl,
```

-68-

B-D-lactosyl,

B-D-maltosyl,

5 β-D-gentiobiosyl,

3-O-B-D-galactopyranosyl- σ -D-arabanopyranosyl or

B-D-maltotriosyl;

or (D):

Q1, Q2, Q4 and Q5 are each methylene;

10

H
$$OR^1$$
 R^1O H R^1O -alkylene(C_2 - C_3)-O H and Q^3 is -C- , -C- ,

ог

and wherein

R1 is

20 B-D-2-acetamido-2-deoxy-glucopyranosyl,

B-D-fucopyranosyl,

B-D-xylopyranosyl,

B-L-xylopyranosyl,

a-L-arabanopyranosyl,

B-D-cellobiosyl,

B-D-gentiobiosyl,

3-O- β -D-galactopyranosyl- α -D-arabanopyranosyl, or

B-maltotriosyl;

or (E):

Q¹, Q², and Q⁵ are each methylene;

Q⁴ is carbonyl or -C-

-69-

H OR1 R10 H

Q3 is -C- or -C-:

5

C₅ is alpha;

C₂₅ is (R); and wherein

R1 is

B-D-galactopyranosyl,

10 β-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl or

B-D-maltotriosyl;

or (F):

15 Q1, Q2, and Q4 are each methylene;

Q⁵ is carbonyl or -C-;

H OR¹ R¹O H

C₅ is alpha;

C₂₅ is (R); and wherein

R1 is

25 B-D-galactopyranosyl,

B-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl or

B-D-maltotriosyl;

with the proviso that $(3\beta,5\alpha,25R)$ -3-[(β -D-cellobiosyl)oxy]spirostane is not included.

2. A compound according to claim 1 wherein Q¹ is carbonyl, -C- or -C-,

- Q^2 , Q^4 and Q^5 are each methylene, Q^3 is $\frac{H}{C_5}$, the C_5 hydrogen is alpha and C_{25} is (R).
- 3. A compound according to claim 2 wherein Q¹ is carbonyl and R¹ is β-D-cellobiosyl.
 - 4. A compound according to claim 2 wherein Q¹ is carbonyl and R¹ is β-D-galactopyranosyl.
 - 5. A compound according to claim 2 wherein Q^1 is carbonyl and R^1 is α -D-cellobiosyl.
- 15
 6. A compound according to claim 2 wherein Q¹ is carbonyl and R¹ is β-D-glucopyranosyl.
 - 7. A compound according to claim 2 wherein Q^1 is carbonyl and R^1 is β -D-lactosyl.
- A compound according to claim 2 wherein Q¹ is carbonyl and R1 is β-D-maltosyl.
 - 9. A compound according to claim 2 wherein Q¹ is carbonyl and R1 is β-D-maltotriosyl.
- 25 10. A compound according to claim 2 wherein Q¹ is -C- and R¹ is β-D-cellobiosyl.
 - 11. A compound according to claim 2 wherein Q¹ is -C- and R¹ is β-D
- cellobiosyl.
 - 12. A compound according to claim 1 wherein Q1, Q4 and Q5 are each methylene,

H OH HO H H OR¹
Q² is
$$^{-C-}$$
 or $^{-C-}$, Q³ is $^{-C-}$, the C₅ hydrogen is alpha and C₂₅ is (R).

- 13. A compound according to claim 12 wherein Q² is -C- and R¹ is β-D-cellobiosyl.
 - 14. A compound according to claim 1 wherein Q¹ is carbonyl, -C- or -C-
- 10 $\frac{H}{Q^2}$ is carbonyl, $\frac{H}{Q^2}$ or $\frac{H}{Q^2}$ or $\frac{H}{Q^2}$ or $\frac{H}{Q^2}$ and $\frac{H}{Q^5}$ are each methylene, $\frac{H}{Q^2}$ is (R), and the $\frac{H}{Q^5}$ hydrogen is alpha.
- 15. A compound according to claim 14 wherein Q¹ is carbonyl, Q² is carbonyl and R¹ is B-D-cellobiosyl.
 - 16. A compound according to claim 14 wherein Q¹ is carbonyl, Q² is -C- and R² is B-D-cellobiosyl.
- 17. A compound according to claim 14 wherein Q¹ is carbonyl, Q² is -C- and R¹ is B-D-lactosyl.
- 18. A compound according to claim 14 wherein Q¹ is -C-, Q² is carbonyl and R¹ is β-D-cellobiosyl.
- 19. A compound according to claim 14 wherein Q¹ is -C-, Q² is carbonyl and R¹ is β-D-cellobiosyl.

- 20. A compound according to claim 1 wherein Q1, Q4 and Q5 are each methylene,
- Q^2 is carbonyl, Q^3 is C^2 , the C_5 hydrogen is alpha and C_{25} is (R).
 - 21. A compound according to claim 20 wherein R¹ is β-D-lactosyl.
- 22. A compound according to claim 20 wherein R¹ is β-D-cellobiosyl.
 - 23. A compound according to claim 1 wherein Q1 and Q2, Q4 and Q5 are each

methylene,
$$Q^3$$
 is C^2 and C_{25} is (R).

15

- 24. A compound according to claim 23 wherein the C₅ hydrogen is beta and R¹ is β-D-cellobiosyl.
- 25. A compound according to claim 23 wherein the C₅ hydrogen is alpha and R¹ is
 B-D-gentiobiosyl.
 - 26. A compound according to claim 1 wherein Q^1 , Q^2 and Q^5 are each

methylene, Q^3 is C^{-1} , Q^4 is carbonyl, the C_5 hydrogen is alpha and C_{25} is (R).

25

- 27. A compound according to claim 26 wherein R¹ is β-D-cellobiosyl.
- 28. A compound according to claim 1 wherein Q^1 , Q^2 and Q^4 are each

methylene, Q3 is -C-, Q5 is carbonyl, the C5 hydrogen is alpha and C25 is (R).

29. A compound according to claim 28 wherein R¹ is β-D-cellobiosyl.

30. A method for controlling hypercholesterolemia or atherosclerosis in a mammal comprising administering to a mammal suffering from hypercholesterolemia or atherosclerosis a hypercholesterolemia or atherosclerosis controlling amount of a Formula I spirostanyl glycoside

10

5

15

Formula I

wherein

either (A):

20

25

30

H
$$OR^1$$
 R^1O H R^1O -alkylene(C_2 - C_3)-O H O^3 is $-C$ -, $-C$ -,

Οľ

$$R^{1}0$$
-alkylene(C_{2} - C_{3})-0 H

Q4 and Q5 are both methylene;

and wherein

5

R1 is

B-D-glucopyranosyl,

B-D-glucopyranuronosyl,

β-D-2-acetamido-2-deoxy-glucopyranosyl,

B-D-galactopyranosyl,

B-D-fucopyranosyl,

B-L-fucopyranosyl,

15 B-D-xylopyranosyl,

B-L-xylopyranosyl,

a-D-arabanopyranosyl,

a-L-arabanopyranosyl,

20 α-D-cellobiosyi,

B-D-cellobiosyl,

B-D-lactosyl,

B-D-maltosyl,

25 β-D-gentiobiosyl,

3-O-B-D-galactopyranosyl-a-D-arabanopyranosyl or

B-D-maltotriosyl;

or (B):

30

Q1, Q2, and Q5 are each methylene;

Q³ is -C- or -C-;

Q⁴ is carbonyl or -C-;

5

C₅ is alpha;

C₂₅ is (R);

and wherein

R¹ is

B-D-galactopyranosyl,

B-D-cellobiosyl,

15 B-D-lactosyl,

B-D-maltosyl or

B-D-maltotriosyl;

or (C):

Q¹, Q² and Q⁴ are each methylene;

25

Q⁵ is carbonyl or -C-;

C_s is alpha;

C₂₅ is (R);

and wherein;

R¹ is

35 B-D-galactopyranosyl,

B-D-cellobiosyl,

B-D-lactosyl,

5

B-D-maltosyi or

B-D-maltotriosyl;

with the proviso that

(38,5a,25R)-3-[(a-D-cellobiosyl)oxy]spirostane,

(3B,5a,25R)-3-[(B-D-glucopyranosyl)oxy]spirostane,

(3B,5a,25R)-3-[(B-D-cellobiosyl)oxy]spirostane or

(3B,5a,25R)-3-[(B-D-galactopyranosyl)oxy]spirostan-12-one are not included.

15

31. The method according to claim 30 wherein $Q^1,\,Q^2,\,Q^4$ and Q^5 are each

methylene, C₂₅ is (R) and Q³ is -C-

- 32. The method according to claim 31 wherein the C_5 hydrogen is beta and R^1 is β -D-cellobiosyl.
- 33. The method according to claim 31 wherein the C_5 hydrogen is alpha and R^1 is β -D-glucopyranuronosyl.
 - 34. The method according to claim 31 wherein the C_5 hydrogen is alpha and R^1 is β -D-maltosyl.
- 35. The method according to claim 31 wherein the C₅ hydrogen is alpha and R¹ is B-D-lactosyl.
 - 36. The method according to claim 31 wherein the $C_{\scriptscriptstyle 5}$ hydrogen is alpha and $R^{\scriptscriptstyle 1}$ is β -D-gentiobiosyl.

5

- 37. The method according to claim 31 wherein the C_5 hydrogen is alpha and R^1 is β -D-galactopyranosyl.
- 38. The method according to claim 30 wherein Q¹ is carbonyl, -C- or -C-,
- 10 Q², Q⁴ and Q⁵ are each methylene, Q³ is -C-, C₂₅ is (R) and the C₅ hydrogen is alpha.
- 39. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is β-D-cellobiosyl.
 - 40. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is β-D-galactopyranosyl.
- 20 41. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is α-D-cellobiosyl.
 - 42. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is β-D-glucopyranosyl.
- 25
 43. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is β-D-maltosyl.
 - 44. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is β-D-maltotriosyl.
 - 45. The method according to claim 38 wherein Q¹ is carbonyl and R¹ is β-D-lactosyl.

- 46. The method according to claim 38 wherein Q' is -C- and R' is β-D-5 cellobiosyl.
- 47. The method according to claim 38 wherein Q¹ is -C- and R¹ is β-Dcellobiosyl.
 - 48. The method according to claim 30 wherein Q1, Q4 and Q5 are each methylene,
- 15 H OH HO H H OR 1 Q² is carbonyl, $^{-C-}$ or $^{-C-}$, Q³ is $^{-C-}$, C₂₅ is (R) and the C₅ hydrogen is

alpha.

- 20 49. The method according to claim 48 wherein Q² is carbonyl and R¹ is β-D-cellobiosyl.
 - 50. The method according to claim 48 wherein Q² is carbonyl and R¹ is β-D-lactosyl.
- 51. The method according to claim 48 wherein Q² is -C- and R¹ is β-D-cellobiosyl.
 - 52. The method according to claim 48 wherein Q² is -C- and R¹ is β-D-galactopyranosyl.

5

53. The method according to claim 30 wherein Q1 is carbonyl, -C- or -C-,

H OH HO H H $0R^{2}$ Q² is carbonyl, $-C^{-}$ or $-C^{-}$, Q³ is $-C^{-}$, Q⁴ and Q⁵ are each methylene, C_{25}

is (R), and the C₅ hydrogen is alpha.

- 54. The method according to claim 53 wherein Q^1 is carbonyl, Q^2 is carbonyl and R^1 is β -D-cellobiosyl.
- 15
 55. The method according to claim 53 wherein Q¹ is carbonyl, Q² is -C- and R¹ is β-D-cellobiosyl.
- 20
 56. The method according to claim 53 wherein Q¹ is carbonyl, Q² is -C- and R is β-D-lactosyl.
- 57. The method according to claim 53 wherein Q¹ is -C- , Q² is carbonyl and R¹ is B-D-cellobiosyl.
- 58. The method according to claim 53 wherein Q^1 is a $-C^-$, Q^2 is carbonyl and

R¹ is B-D-cellobiosyl.

59. The method according to claim 30 wherein Q^1 , Q^2 and Q^5 are each

5 methylene, Q^3 is C^{-1} , Q^4 is carbonyl, the C_5 hydrogen is alpha and C_{25} is (R).

- 60. The method according to claim 59 wherein R¹ is β-D-cellobiosyl.
- 61. The method according to claim 30 wherein Q^1 , Q^2 and Q^4 are each 10 H $_{10}^{0}$ $_{10}^{1}$ $_{10}^$
- 15 62. The method according to claim 61 wherein R¹ is β-D-cellobiosyl.
 - 63. The pharmaceutical composition for the control of hypercholesterolemia or atherosclerosis in mammals which comprises a compound of claim 1 and a pharmaceutically acceptable carrier.
- 20 64. The composition comprising a hydrate of a compound according to claim 1.

25

International Application No

L CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all)6									
	Patent Classification (IPC) or to both Nation								
II. FIELDS SEARCHED									
•	. Minimum Documentation Searched?								
Classification System		Classification Symbols							
Int.C1. 5	C07J ; A61K		·						
	Documentation Searched to the Extent that such Docum	other than Minimum Documentation nents are Included in the Fields Searched ⁸							
	IDERED TO BE RELEVANT		Coin No.13						
Category Citatio	n of Document, 11 with indication, where ap	propriate, of the relevant passages 12	Relevant to Claim No.13						
20 c abs G.	MICAL ABSTRACTS, vol. 10 July 1987, Columbus, Ohi tract no. 22075, S. SIDHU ET AL 'Effects Tigogenin-cellobioside	of Soy Saponins	1,63						
Upt Glud page see & N vol	ake of Cholesterol, Choleose' 499 ;column 1 ; abstract JTR. REP. INT 35, no. 3, 1987, es 615 - 623	late and							
		-/	-						
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filling date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filling date but later than the priority date claimed "A" document member of the same patent family "A" document member of the same patent family		the application but bry underlying the aimed invention acconsidered to aimed invention noise step when the sother such docu- to a person skilled							
IV. CERTIFICATION	cian of the International Court	Date of Mailing of this International So	earch Report						
1	etion of the International Search AUGUST 1993	1 5. 09. 93							
International Searching Authority EUROPEAN PATENT OFFICE		Signature of Authorized Officer WATCHORN P.W.							

III. DOCUM	IENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)	
Category °	Citation of Document, with Indication, where appropriate, of the relevant passages	Relevant to Claim No.
X	STEROIDS: STRUCTURE, FUNCTION, AND	1,63
	REGULATION vol. 48, no. 3-4, 1986, STONEHAM, MA US pages 197 - 211 M. R. MALINOW ET AL 'Effects of alpha- and beta-Tigogenin Cellobiosides on	
	Cholesterol Absorption' see the whole document	
X	EP,A,O 159 431 (MEDICAL RESEARCH FOUNDATION OF OREGON) 30 October 1985 see examples 4,5	1,63
X	EP,A,O 403 150 (PFIZER INC.) 19 December 1990 see page 2	1,63
X	ANNALS OF THE NEW YORK ACADEMY OF SCIENCES (ATHEROSCLEROSIS) vol. 454, 1985, NEW YORK, US pages 23 - 27 M. R. MALINOW 'Effects of Synthetic Glycosides on Cholesterol Absorption' see the whole document	1,63
X	US,A,4 461 762 (M. R. MALINOW) 24 July 1984 see column 3 - column 10	1,63
X	JOURNAL OF PHARMACEUTICAL SCIENCES vol. 67, no. 11, 1978, WASHINGTON D.C., US pages 1589 - 1592 R. SEGAL ET AL 'Hemolytic Properties of Synthetic Glycosides' see the whole document	1
X	JUSTUS LIEBIGS ANNALEN DER CHEMIE vol. 699, 1966, WEINHEIM DE pages 212 - 222 R. TSCHESCHE ET AL 'Über Parillin, ein Saponin mit stark verzweigter Zuckerkette' see page 218, line 28 - line 30 see page 216	1

INTERNATIONAL SEARCH REPORT

PCT/US 93/04092

Box I	Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)
This inte	ernational search report has not been established in respect of œrtain claims under Article 17(2)(a) for the following reasons:
1. X	Claims Nos.: because they relate to subject matter not required to be searched by this Authority, namely: Remark: Although claims 30-62 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compounds.
2.	Claims Nos.: because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3.	Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).
Box II	Observations where unity of invention is lacking (Continuation of item 2 of first sheet)
This Inte	ernational Searching Authority found multiple inventions in this international application, as follows:
1.	As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2.	As all searchable claims could be searches without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.	As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4.	No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Remark	on Protest The additional search fees were accompanied by the applicant's protest. No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.

9304092 US 74430 SA

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 31/08/93

Patent document cited in search report	Publication date 30-10-85	Patent family member(s)		Publication date
EP-A-0159431		US-A- AU-B- AU-A- CA-A- JP-B- JP-A-	4602005 580005 3097784 1246544 4056840 60224697	22-07-86 22-12-88 24-10-85 13-12-88 09-09-92 09-11-85
EP-A-0403150	19-12-90	US-A- AU-B- AU-A- CA-A- JP-A-	5010185 619986 5707990 2018722 3031297	23-04-91 06-02-92 01-08-91 13-12-90 12-02-91
US-A-4461762	24-07-84	US-A- US-A-	4602003 4602005	22-07-86 22-07-86

This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:			
BLACK BORDERS			
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES			
☐ FADED TEXT OR DRAWING			
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING			
☐ SKEWED/SLANTED IMAGES			
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS			
GRAY SCALE DOCUMENTS			
LINES OR MARKS ON ORIGINAL DOCUMENT			
REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY			
□ OTHER:			

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.